



FRAC CODE LIST 2: Fungicides sorted by modes of action

INTRODUCTION

The following table lists commercial fungicides according to their mode of action and resistance risk. The most important bactericides are also included.

The Table headings are defined as:

MOA Code

Different letters (A to I, with added numbers) were used to distinguish fungicide groups according to their general mode of action (MOA) in the biosynthetic processes of fungi. The grouping was made according to the processes in primary metabolism from nucleic acids synthesis (A) to secondary metabolism with melanin synthesis (I) at the end of the list, followed by host plant defence inducers (P), recent molecules with an unknown mode of action and unknown resistance risk (U, transient status, mostly not longer than 8 years, until information about mode of action and mechanism of resistance becomes available), and multi-site inhibitors (M).

Target Site of Action

If available the biochemical mode of action is given. In many cases the precise target site is not known. However, a grouping can be made due to cross resistance profiles within a group or in relation to other groups.

Group Name

The Group Names listed are widely accepted in literature. They are based on different sources (mode of action, first important representative, chemical group).

Chemical Group

Sub-grouping due to chemical considerations.

Common name

Accepted (or proposed) common name for an individual active ingredient expected to appear on the product label as definition of the product.

Comments on Resistance

If field resistance is known to one member of the Group, it is most likely but not exclusively valid that cross resistance to other Group members will be present. There is increasing evidence that cross resistance may not be clearly visible between Group members and that the degree of the effect can differ both between group members and fungal species or even within species. For the latest information on resistance and cross resistance status of a particular fungus-fungicide complex, you are

advised to contact your local FRAC representative, product manufacturer's representative or crop protection advisor. The intrinsic risk for resistance evolution to a given fungicide group is estimated to be **low, medium or high** according to the principles described in FRAC Monographs 1, 2 and 3. Resistance management is driven by pathogen risk and agronomic risk (see FRAC pathogen risk list).

Similar classification lists of fungicides have been published by T. Locke on behalf of FRAG – UK (Fungicide Resistance, August 2001), and by P. Leroux (Classification des fongicides agricoles et résistance, Phytoma, La Défense des Végétaux, No. 554, 43-51, November 2002).

FRAC Code

Numbers and letters are used to distinguish the fungicide groups according to their cross resistance behaviour. The numbers were assigned primarily according to the time of product introduction to the market. The letters refer to P = host plant defence inducers, M = multi-site inhibitors, and U = unknown mode of action and unknown resistance risk.

Last update: December 2006

Next update: December 2007

MOA	TARGET SITE AND CODE	GROUP NAME	CHEMICAL GROUP	COMMON NAME	COMMENTS	FRAC CODE
A: Nucleic acids synthesis	A1: RNA polymerase I	PA – fungicides (PhenylAmides)	acylalanines	benalaxyl furalaxyl metalaxyl metalaxyl-M (=mefenoxam)	Resistance and cross resistance well known in various Oomycetes but mechanism unknown. High risk. See FRAC Phenylamide Guidelines for resistance management	4
			oxazolidinones	oxadixyl		
			butyrolactones	ofurace		
	A2: adenosin-deaminase	hydroxy-(2-amino-) pyrimidines	hydroxy-(2-amino-) pyrimidines	bupirimate dimethirimol ethirimol	Medium risk Resistance and cross resistance known in powdery mildews. Resistance management required.	8
	A3: DNA/RNA synthesis (proposed)	heteroaromatics	isoxazoles	hymexazole	Resistance not known	32
			isothiazolones	octhilinone		
A4: DNA topoisomerase type II (gyrase)	carboxylic acids	carboxylic acids	oxolinic acid	Bactericide. Resistance known. Risk unknown. Resistance management required.	31	
B: Mitosis and cell division	B1: β -tubuline assembly in mitosis	MBC - fungicides (Methyl Benzimidazole Carbamates)	benzimidazoles	benomyl carbendazim fuberidazole thiabendazole	Resistance common in many fungal species. Several target site mutations, mostly E198A/G/K, F200Y in β -tubulin gene Positive cross resistance between the group members. Negative cross resistance to N-Phenylcarbamates High risk. See FRAC Benzimidazole Guidelines for resistance management.	1
			thiophanates	thiophanate thiophanate-methyl		
	B2: β -tubulin assembly in mitosis	N-phenyl carbamates	N-phenyl carbamates	diethofencarb	Resistance known. Target site mutation E198K. Negative cross resistance to benzimidazoles. High risk. Resistance management required.	10
	B3: β -tubulin assembly in mitosis	benzamides	toluamides	zoxamide	Low to medium risk. Resistance management required.	22
	B4: cell division (proposed)	phenylureas	phenylureas	pencycuron	Resistance not known	20
	B5: delocalisation of spectrin-like proteins	benzamides	acylpicolides	fluopicolide	Resistance not known	43

MOA	TARGET SITE AND CODE	GROUP NAME	CHEMICAL GROUP	COMMON NAME	COMMENTS	FRAC CODE
C. Respiration	C1: complex I NADH Oxidoreductase	pyrimidinamines	pyrimidinamines	diflometorim	Resistance not known	39
	C2: complex II: succinate-dehydrogenase	carboxamides	benzamides	benodanil flutolanil mepronil	Resistance known for specific fungi. Target site mutation H257L. Medium risk. Resistance management required if used for risky pathogens.	7
			furan carboxamides	fenfuram		
			oxathiin carboxamides	carboxin oxycarboxin		
			thiazole carboxamides	thifluzamide		
			pyrazole carboxamides	furametpyr penthiopyrad		
	C3: complex III: cytochrome bc1 (ubiquinol oxidase) at Qo site (<i>cyt b gene</i>)	QoI-fungicides (Quinone outside Inhibitors)	methoxy-acrylates	azoxystrobin enestrobin picoxystrobin	Resistance known in various fungal species. Target site mutations G143A, F129L and additional mechanisms.	11
			methoxy-carbamates	pyraclostrobin		
			oximino acetates	kresoxim-methyl trifloxystrobin	Cross resistance shown between all members of the QoI group.	
			oximino-acetamides	dimoxystrobin metominostrobin orysastrobin		
			oxazolidinediones	famoxadone		
			dihydro-dioxazines	fluoxastrobin	High risk. See FRAC QoI Guidelines for resistance management.	
			imidazolinones	fenamidone		
benzyl-carbamates	pyribencarb					
C4: complex III: cytochrome bc1(ubiquinol-oxidase)	Qil - fungicides (Quinone inside Inhibitors)	cyano-imidazole	cyazofamid	Resistance risk unknown but assumed to be medium to high (mutations at target site known in model organisms). Resistance management required.	21	
		sulfamoyl-triazole	amisulbrom			
C5: uncouplers of oxidative phosphorylation		dinitrophenyl crotonates	binapacryl dinocap	Resistance not known	29	
		2,6-dinitro-anilines	fluazinam	Low risk. However, resistance claimed in Botrytis in Japan.		
		pyrimidinone-hydrazones	ferimzone	Resistance not known		
C6: inhibitors of oxidative phosphorylation, ATP synthase	organo tin compounds	tri phenyl tin compounds	fentin acetate fentin chloride fentin hydroxide	Some resistance cases known. Low to medium risk.	30	
C7: ATP production (proposed)	thiophene-carboxamides	thiophene-carboxamides	silthiofam	Resistance reported. Risk low	38	

MOA	TARGET SITE AND CODE	GROUP NAME	CHEMICAL GROUP	COMMON NAME	COMMENTS	FRAC CODE
D: Amino acids and protein synthesis	D1: methionine biosynthesis (proposed) (<i>cgs gene</i>)	AP - fungicides (Anilino-Pyrimidines)	anilino-pyrimidines	cyprodinil mepanipyrim pyrimethanil	Resistance known in Botrytis and Venturia, sporadically in Oculimacula . Medium risk. See FRAC Anilinopyrimidine Guidelines for resistance management.	9
	D2: protein synthesis	enopyranuronic acid antibiotic	enopyranuronic acid antibiotic	blastidicin-S	Low to medium risk. Resistance management required.	23
	D3: protein synthesis	hexopyranosyl antibiotic	hexopyranosyl antibiotic	kasugamycin	Resistance known. Medium risk. Resistance management required.	24
	D4: protein synthesis	glucopyranosyl antibiotic	glucopyranosyl antibiotic	streptomycin	Bactericide. Resistance known. High risk. Resistance management required.	25
	D5: protein synthesis	tetracycline antibiotic	tetracycline antibiotic	oxytetracycline	Bactericide. Resistance known. High risk. Resistance management required.	41
E: Signal transduction	E1: G-proteins in early cell signalling (proposed)	quinolines	quinolines	quinoxifen	Resistance known. Medium risk. Resistance management required.	13
	E2: MAP/Histidine-Kinase in osmotic signal transduction (<i>os-2, HOG1</i>)	PP-fungicides (PhenylPyrroles)	phenylpyrroles	fenpiclonil fludioxonil	Resistance found sporadically, mechanism speculative. Low to medium risk. Resistance management required.	12
	E3: MAP/Histidine-Kinase in osmotic signal transduction (<i>os-1, Daf1</i>)	dicarboximides	dicarboximides	chlozolinate iprodione procymidone vinclozolin	Resistance common in Botrytis and some other pathogens. Several mutations in OS-1, mostly I365S Cross resistance common between the group members. Medium to high risk. See FRAC Dicarboximide Guidelines for resistance management.	2

MOA	TARGET SITE AND CODE	GROUP NAME	CHEMICAL GROUP	COMMON NAME	COMMENTS	FRAC CODE
F: Lipids and membrane synthesis	F1	formerly dicarboximides				
	F2: phospholipid biosynthesis, methyltransferase	phosphorothiolates	phosphorothiolates	edifenphos iprobenfos (IBP) pyrazophos	Resistance known for specific fungi. Low to medium risk. Resistance management required if used for risky pathogens.	6
		dithiolanes	dithiolanes	isoprothiolane		
	F3: lipid peroxidation (proposed)	AH-fungicides (Aromatic Hydrocarbons) (chlorophenyls, nitroanilines)	aromatic hydrocarbons	biphenyl chloroneb dicloran quintozene (PCNB) tecnazene (TCNB) tolclofos-methyl	Resistance known to some fungi. Low to medium risk. Cross resistance patterns complex due to different activity spectra.	14
		heteroaromatics	1,2,4-thiadiazoles	etridiazole		
	F4: cell membrane permeability, fatty acids (proposed)	carbamates	carbamates	iodocarb propamocarb prothiocarb	Low to medium risk. Resistance management required.	28
	F5: phospholipid biosynthesis and cell wall deposition (proposed)	CAA-fungicides (Carboxylic Acid Amides)	cinnamic acid amides	dimethomorph flumorph	Resistance known in <i>Plasmopara viticola</i> but not in <i>Phytophthora infestans</i> . Cross resistance between all members of the CAA group. Low to medium risk. See FRAC CAA Guidelines for resistance management	40
			valinamide carbamates	benthiavalicarb iprovalicarb valiphenal		
			mandelic acid amides	mandipropamid		

MOA	TARGET SITE AND CODE	GROUP NAME	CHEMICAL GROUP	COMMON NAME	COMMENTS	FRAC CODE
G: Sterol biosynthesis in membranes	G1: C14-demethylase in sterol biosynthesis (<i>erg11/cyp51</i>)	DMI-fungicides (DeMethylation Inhibitors) (SBI: Class I)	piperazines	triforine	There are great differences in the activity spectra of the different DMI fungicides.	3
			pyridines	pyrifenoxy		
			pyrimidines	fenarimol nuarimol		
			imidazoles	imazalil oxpoconazole pefurazoate prochloraz triflumizole		
			triazoles	azaconazole bitertanol bromuconazole cyproconazole difenoconazole diniconazole epoxiconazole fenbuconazole fluquinconazole flusilazole flutriafol hexaconazole imibenconazole ipconazole metconazole myclobutanil penconazole propiconazole prothioconazole simeconazole tebuconazole tetraconazole triadimefon triadimenol triticonazole		
	G2: Δ^{14} -reductase and $\Delta^8 \rightarrow \Delta^7$ -isomerase in sterol biosynthesis (<i>erg24, erg2</i>)	Amines ("Morpholines") (SBI: Class II)	morpholines	aldimorph dodemorph fenpropimorph tridemorph	Decreased sensitivity for powdery mildews. Cross resistance within the group generally found but not to other SBI classes. Low to medium risk. See FRAC SBI Guidelines for resistance management.	5
			piperidines	fenpropidin piperalin		
			spiroketal-amines	spiroxamine		
	G3: 3-keto reductase, C4-demethylation (<i>erg27</i>)	hydroxyanilides (SBI: Class III)	hydroxyanilides	fenhexamid	Low to medium risk. Resistance management required.	17
	G4: squalene-epoxidase in sterol biosynthesis (<i>erg1</i>)	(SBI class IV)	thiocarbamates	pyributicarb	Resistance not known, fungicidal and herbicidal activity	18
			allylamines	naftifine terbinafine	Medical fungicides only	

MOA	TARGET SITE AND CODE	GROUP NAME	CHEMICAL GROUP	COMMON NAME	COMMENTS	FRAC CODE
H: Glucan synthesis	H3: trehalase and inositol-biosynthesis	glucopyranosyl antibiotic	glucopyranosyl antibiotic	validamycin	Resistance not known	26
	H4: chitin synthase	polyoxins	peptidyl pyrimidine nucleoside	polyoxin	Resistance known. Medium risk. Resistance management required.	19
I: Melanin synthesis in cell wall	I1: reductase in melanin biosynthesis	MBI-R (Melanin Biosynthesis Inhibitors – Reductase)	isobenzofuranone	fthalide	Resistance not known	16.1
			pyrrolo-quinolinone	pyroquilon		
			triazolobenzothiazole	tricyclazole		
	I2: dehydratase in melanin biosynthesis	MBI-D (Melanin Biosynthesis Inhibitors – Dehydratase)	cyclopropane-carboxamide	carpropamid	Resistance known. Medium risk. Resistance management required.	16.2
			carboxamide	diclocymet		
			propionamide	fenoxanil		
P: Host plant defence induction	P1: salicylic acid pathway	benzo-thiadiazole BTH	benzo-thiadiazole BTH	acibenzolar-S-methyl	Resistance not known	P
	P2	benzothiazole	benzothiazole	probenazole (also antibacterial and antifungal activity)	Resistance not known	
	P3	thiadiazole-carboxamide	thiadiazole-carboxamide	tiadinil isotianil	Resistance unknown	

MOA	TARGET SITE AND CODE	GROUP NAME	CHEMICAL GROUP	COMMON NAME	COMMENTS	FRAC CODE
Unknown mode of action (U numbers not appearing in the list derive from reclassified fungicides)	unknown	cyanoacetamide-oxime	cyanoacetamide-oxime	cymoxanil	Resistance claims described. Low to medium risk. Resistance management required.	27
	unknown	phosphonates	ethyl phosphonates	fosetyl-Al	Few resistance cases reported in few pathogens. Low risk	33
				phosphorous acid and salts		
	unknown	phthalamic acids	phthalamic acids	teclofthalam (Bactericide)	Resistance not known	34
	unknown	benzotriazines	benzotriazines	triazoxide	Resistance not known	35
	unknown	benzene-sulfonamides	benzene-sulfonamides	flusulfamide	Resistance not known	36
	unknown	pyridazinones	pyridazinones	diclomezine	Resistance not known	37
	unknown	thiocarbamate	thiocarbamate	methasulfocarb	Resistance not known	42
	microtubule disruption (proposed)	thiazole carboxamide	thiazole carboxamide	ethaboxam	Resistance not known	U5
	unknown	phenyl-acetamide	phenyl-acetamide	cyflufenamid	Resistance in Sphaerotheca Resistance management required	U6
	unknown	quinazolinone	quinazolinone	proquinazid	Resistance not known	U7
	unknown	benzophenone	benzophenone	metrafenone	Resistance not known	U8
	unknown					U10
Not classified	unknown	diverse	diverse	mineral oils, organic oils, potassium bicarbonate, material of biological origin	Resistance not known	NC

MOA	TARGET SITE AND CODE	GROUP NAME	CHEMICAL GROUP	COMMON NAME	COMMENTS	FRAC CODE
Multi-site contact activity	multi-site contact activity	inorganic	inorganic	copper (different salts)	Generally considered as a low risk group without any signs of resistance developing to the fungicides * For dodine, resistance was reported in <i>Venturia inaequalis</i> suggesting that dodine may not be a multi-site inhibitor. Resistance management recommended No cross resistance between group members M1 to M9	M1
		inorganic	inorganic	sulphur		M2
		dithiocarbamates and relatives	dithio-carbamates and relatives	ferbam mancozeb maneb metiram propineb thiram zineb ziram		M3
		phthalimides	phthalimides	captan captafol folpet		M4
		chloronitriles (phthalonitriles)	chloronitriles (phthalonitriles)	chlorothalonil		M5
		sulfamides	sulfamides	dichlofluanid tolylfluanid		M6
		guanidines	guanidines	dodine* guazatine iminocladine		M7
		triazines	triazines	anilazine		M8
		quinones (anthraquinones)	quinones (anthraquinones)	dithianon		M9