

Mode of action of CAAs in Oomycetes

- Albert, G., Thomas, A. and Guehne, M. (1991): Fungicidal activity of dimethomorph on different stages in the life cycle of *Phytophthora infestans* and *Plasmopara viticola*. ANPP – 3rd International Conference on Plant Diseases, Bordeaux, pp. 887-894
- Cohen, Y. and Gisi, U. (2007): Differential activity of carboxylic acid amide fungicides against various development stages of *Phytophthora infestans*. *Phytopathology* 97, 1274-1283
- Cohen, Y., Baider, A. and Cohen, BH. (1995): Dimethomorph activity against oomycete fungal plant pathogens. *Phytopathology* 85, 1500-1506
- Cohen Y., Rubin, E. and Gotlieb, D. (2008): Activity of carboxylic acid amide (CAA) fungicides against *Bremia lactucae*. *European Journal of Plant Pathology* 122:169–183,
- Gisi, U., Lamberth, C., Mehl, A. and Seitz, T. (2007). Carboxylic Acid Amide (CAA) fungicides. In W. Krämer and U. Schirmer, eds., *Modern Crop Protection Compounds* (pp. 651-674), Wiley-VCH, Weinheim, Germany
- Gisi, U. and Sierotzki, H. (2008): Fungicide modes of action and resistance in downy mildews. *European Journal of Plant Pathology* 122, 157-167
- Griffiths, R.G., Dancer, J., O'Neill, E. and Harwood, J.L. (2003): A mandelamide pesticide alters lipid metabolism in *Phytophthora infestans*. *New Phytologist* 158, 345-353
- Mehl, A. and Buchenauer, H. (2002): Investigations of the biochemical mode of action of iprovalicarb. In: HW Dehne, U Gisi, KH Kuck, PE Russell, H Lyr, eds., *Modern Fungicides and Antifungal Compounds III*, AgroConcept, Bonn, Germany, pp. 75-82
- Zhu, S.S., Liu, X.L., Liu, P.F., Li, J.Q., Wang, H.M., Yuan, S.K. and Si, N.G. (2007). Flumorph is a novel fungicide that disrupts microfilament organization in *Phytophthora melonis*. *Phytopathology* 97, 643-649