

Anilinopyrimidines  
Working Group

**FRAC**  
FUNGICIDE RESISTANCE  
ACTION COMMITTEE

## **Anilinopyrimidines (AP's) Working Group**

**Meeting on December 13, 2016, 8:30 am - 12:30 am  
Protocol of the discussions and use recommendations of the AP's Working  
Group of the Fungicide Resistance Action Committee (FRAC)**

---

### **Participants**

BASF	Randall Gold Gerd Stammler
Bayer	<u>Andreas Mehl</u> Sophie Huvier-Boutin
K-I Chemical	Yu Aoki Takumi Katsumata
Syngenta	Stefano Torriani Birgit Forster

### **Venue:**

Lindner Main Plaza Hotel, Frankfurt, Germany

---

**ANTI-TRUST GUIDELINES (FROM FRAC CONSTITUTION) WERE SHOWN AT  
THE START OF THE MEETING**

Source: [www.frac.info](http://www.frac.info)  
Dec 2016

## **1. Monitoring Results 2016 (FRAC members)**

### **1.1 *Botrytis* results**

#### Vineyards

(BASF, Bayer, K-I Chemical, Syngenta)

Extensive monitoring studies in Europe, Chile, and New Zealand have been carried out for more than a decade by Bayer, K-I Chemical, Syngenta, and BASF. In 2016, sensitivity data from commercial vineyards and trial sites were presented for France, Germany, Austria, Chile, New Zealand, Croatia, Italy, Greece, Czech Republic, Spain, Romania, Hungary, Portugal, Slovakia, and Switzerland.

Data from these studies show that frequencies of resistant strains varied from zero to moderate with regional variability. These results are comparable to the observations made during the last ten years.

Products, applied according to the FRAC-AP guidelines in grape spray programs, maintained very good performance in the field.

#### Strawberries

(BASF, Syngenta)

Sensitivity monitoring was carried out during 2016 in Belgium, Czech Republic, Denmark, France, Germany, Greece, Hungary, Italy, Latvia, Netherlands, Poland, Slovenia, Spain, and the UK from commercial locations and trial sites.

The data show that the frequency of resistant isolates is moderate, fluctuating from field to field, and ranging from zero to high. Compared to the last ten years, the frequency of resistant isolates in the monitored populations remained also stable. Products, applied according to the FRAC-AP guidelines in strawberry spray programs, provided good control in commercial situations.

#### Vegetables

(Syngenta)

Sensitivity monitoring in tomato, peas, and beans was carried out during 2016 in the Netherlands, Belgium, Czech Republic, Italy, Serbia, France, and Poland with a limited number of samples obtained from commercial locations and trial sites. Overall, the data show a low frequency of resistant isolates in tomato and beans.

Evidence from field and laboratory trials has shown that there is a medium resistance risk of *Botrytis* to APs. Good agronomic practices and strict adherence to the FRAC AP use guidelines are crucial to ensure that APs remain effective due to the risk of increasing occurrence of multiple resistant strains and multidrug resistant (MDR) strains, particularly in soft fruits. For more information on these different resistance types see quick references.

## **1.2. *Venturia* results**

(BASF, Syngenta)

Monitoring studies for 2016 were presented.

Samples from France, Germany, Italy, Croatia, Czech Republic, Romania, Belgium, Hungary, Poland, Portugal, Bulgaria, Latvia, Lithuania, Spain, Switzerland, Turkey, Greece, and the UK were analyzed. Based on dose-response to APs using *in vivo* biotests, populations are classified as either sensitive, moderately adapted or resistant.

In 2016, the trend for lower resistance frequencies at commercial and trial site locations, as already observed in 2015, was confirmed. This was linked to the detection of more sensitive and less moderately adapted populations in all apple growing areas monitored.

---

## **2. Use Recommendations**

The purpose of the use guidelines for AP containing products is to maintain the sensitivity in the target pathogens and to prevent crop losses due to resistant pathogen populations.

### **2.1 General AP's Guidelines (all crops)**

Where different AP-containing products are used in one season, the cumulative number of applications with cyprodinil-, pyrimethanil- or mepanipyrim-containing products must not exceed the maxima as mentioned below.

For sound resistance management, good agricultural practices, including phytosanitary measures and crop protection, should be followed not only in commercial practice, but also in nurseries.

The use recommendations were reviewed during the meeting on December 13<sup>th</sup>, 2016. The *Botrytis* and *Venturia* guidelines have not been changed.

### **2.2 *Botrytis* Guidelines**

- Where up to three treatments are made per season, the number of applications of AP-containing products is limited to one.
- In situations where four to six *Botrytis* treatments are made per crop and season, a maximum of two applications with AP-containing products are recommended.
- In specific situations where seven or more *Botrytis* treatments are required per crop and season, a maximum of three applications with AP-containing products is recommended and not more than two consecutive applications.

Source: [www.frac.info](http://www.frac.info)  
Dec 2016

- For specific crops and products, follow use recommendations of individual companies.

### **2.3 *Venturia* Guidelines**

- Apply a maximum of four AP-containing products per season.
  - In locations where moderately adapted or resistant populations have been reported, use APs only in mixture with an effective non cross resistant scab fungicide.
  - Individual products should always be used at recommended dose rates and during the period when they are most effective.
  - Curative use only in conjunction with reliable scab warning systems.
- 

The next AP FRAC Working Group meeting is scheduled for December, the 12<sup>th</sup>, 2017.