



## **Anilinopyrimidines (AP's) Working Group**

**Meeting on January 17, 2022, 1:00 pm - 5:00 pm**  
**Protocol of the discussions and use recommendations of the AP's Working Group**  
**of the Fungicide Resistance Action Committee (FRAC)**

---

### **Participants**

BASF	Nadine Riediger Gerd Stammler
Bayer	<u>Andreas Mehl</u> Daniela Portz
K-I Chemical	Akihiro Moriwaki
Syngenta	Stefano Torriani Thierry Querol

### **Venue:**

virtual on-line Teams-meeting

---

## **Disclaimer**

The technical information contained in the global guidelines/the website/the publication/the minutes is provided to CropLife International/RAC members, non-members, the scientific community, and a broader public audience.

While CropLife International and the RACs make every effort to present accurate and reliable information in the guidelines, CropLife International and the RACs do not guarantee the accuracy, completeness, efficacy, timeliness, or correct sequencing of such information. CropLife International and the RACs assume no responsibility for consequences resulting from the use of their information, or in any respect for the content of such information, including but not limited to errors or omissions, the accuracy or reasonableness of factual or scientific assumptions, studies, or conclusions.

Inclusion of active ingredients and products on the RAC Code Lists is based on scientific evaluation of their modes of action; it does not provide any kind of testimonial for the use of a product or a judgment on efficacy. CropLife International and the RACs are not responsible for, and expressly disclaim all liability for, damages of any kind arising out of use, reference to, or reliance on information provided in the guidelines.

Listing of chemical classes or modes of action in any of the CropLife International/RAC recommendations must not be interpreted as approval for use of a compound in a given country. Prior to implementation, each user must determine the current registration status in the country of use and strictly adhere to the uses and instructions approved in that country.

Anti-Trust Guidelines (from FRAC Constitution) were shown before meetings started

## **1.1 Botrytis results**

### **Vineyards**

(BASF, Bayer, K-I Chemical, Syngenta)

As in 2020, in 2021, sensitivity data from commercial vineyards and trial sites were presented for France, Italy, and Spain in the major wine growing regions, as well as from Croatia, Germany, Hungary, Poland, Portugal, and Switzerland. Data from these studies show that frequencies of resistant strains was generally low and still varied in Europe from zero to moderate with regional variability.

These results still reflect a stable situation and are comparable to the observations made since more than a decade.

In 2019, sensitivity data from commercial vineyards and trial sites were presented for France, Germany, Austria, Croatia, Italy, Portugal, Czech Republic, Spain, Switzerland, Slovenia, and Hungary.

Frequencies of resistant strains varied in Europe from zero to moderate and in Chile (2018 data) from zero to high especially at trial sites, with regional variability.

Products, applied according to the FRAC-AP guidelines in grape spray programs, maintained good performance in the field.

### **Strawberries**

(BASF, Bayer, Syngenta)

As in 2020, in 2021, sensitivity monitoring was carried out in Belgium, Croatia, Finland, France, Germany, Italy, Poland, Slovenia, and Spain, from commercial locations and trial sites.

Data show that the frequency of resistant isolates is moderate, fluctuating from field to field, and still ranging from zero to high. Compared to data obtained a decade ago, the frequency of resistant isolates in the monitored populations remained stable.

In 2019, sensitivity monitoring was carried out in Croatia, France, Germany, Poland, Portugal, Greece, UK, Netherlands, Slovenia, and Spain, from commercial locations and trial sites.

The frequency of resistant isolates was again moderate, fluctuating from field to field, and ranging from zero to high.

Monitoring studies in 2017 and 2018 with samples originating from 8 Chinese provinces show mostly full sensitive isolates and only single resistant strains.

Products, applied according to the FRAC-AP guidelines in strawberry spray programs, provided good control in commercial situations.

## **Vegetables**

(Syngenta)

As in the last years, in 2020 and 2021, limited sensitivity data for tomato, lettuce, and cucumber showed a variable situation with sensitive and resistant isolates in samples originating from France and Spain.

In 2019, limited sensitivity data for tomato samples collected in a single location in Southern Italy indicated presence of resistant isolates.

In 2017 and 2018, limited sensitivity data for tomato samples collected in Belgium and the Netherlands showed no to moderate frequency of resistant isolates. Limited monitoring was also carried out in beans: one Spanish sample from 2018 showed resistance, while all samples from France were sensitive.

In 2016, European sensitivity monitoring in tomato, peas, and beans was carried out in the Netherlands, Belgium, Czech Republic, Italy, Serbia, France, and Poland with a limited number of samples obtained from commercial locations and trial sites.

Overall, the data show a low frequency of resistant isolates in tomato and beans.

### Monitoring outside of Europe:

In 2018, samples originating from up to 6 Chinese provinces didn't show occurrence of resistant isolates in tomato and cucumber.

In 2017, first monitoring studies with limited number of tomato and cucumber samples from China indicated, similar to the situation in strawberries, besides sensitive isolates also presence of resistant ones.

Evidence from field and laboratory trials has shown that there is a medium resistance risk of *Botrytis* to APs. Good agronomic practices and strict adherence to the FRAC AP use guidelines are crucial to ensure that APs remain effective due to the risk of increasing occurrence of multiple resistant strains and isolates with unspecific, increased efflux activity (often called 'multidrug resistant', MDR), particularly in soft fruits.

For more information on the different AP resistance types see the latest publications on AP related resistance research (Mosbach *et al.*, 2017).

## **1.2. Venturia results**

(BASF, Syngenta)

2021 monitoring is still ongoing.

In 2019 and 2020, samples from Austria, France, Germany, Italy, Belgium, Bulgaria, Hungary, Poland, Portugal, Spain, Greece, Latvia, Croatia, Netherlands, Ukraine, Slovenia, Switzerland, and the UK were analysed. Based on dose-response to APs using *in vivo* biotests, populations are classified as either sensitive, moderately adapted, or resistant.

As in the past years, the vast majority of all studied samples was sensitive, indicating that current resistance management strategies are efficient.

History:

In 2016 and 2017, the trend for lower resistance frequencies at commercial and trial site locations, as already observed in 2015, was confirmed. This was linked to the detection of more sensitive and less moderately adapted populations in all apple growing areas monitored.

### **1.3. *Stemphylium vesicarium***

(Syngenta)

Monitoring data has been presented for isolates originating from Portugal, UK, Croatia, Belgium, Germany, Spain, Greece, Italy, Netherlands, and Poland during 2015-2021, sampled in asparagus and pears.

In summary, the majority of all studied isolates were fully sensitive. However, single resistant isolates were detected in both crops.

### **1.4. *Alternaria solani*** (potatoes)

(Bayer)

2021 monitoring is still ongoing.

In 2020, samples from the Netherlands, Austria, Germany, Norway, Denmark, Sweden, and Belgium were analysed. High sensitivity was reported.

First monitoring data has been generated in 2019 for European isolates originating from Belgium, Germany, Netherlands, Denmark, and Sweden.

All isolates showed high sensitivity in a narrow range.

### **1.5. *Monilinia spp.*** (stone- and pome fruit)

(BASF)

In 2021, samples from Switzerland, Czech Republic, Germany, Spain, France, Greece, Hungary, Italy, Poland, Portugal, and Romania were analysed. AP sensitivity was, overall, high with some variations in the baseline, dependent on the species (*M. laxa*, *M. fructicola*, *M. fructigena*).

First and preliminary monitoring data have been generated in 2020 for European isolates originating from Portugal and Greece.

All isolates were sensitive and EC<sub>50</sub> values showed a normal frequency distribution.

## **2. AP - Use Recommendations**

The purpose of the use guidelines for AP containing products is to maintain the sensitivity in the target pathogens and to prevent crop losses due to resistant pathogen populations.

### **2.1 AP - Guidelines**

For grey mold and apple scab control, specific guidelines have been developed. In general, where different AP-containing products are used in one season, the cumulative number of applications with cyprodinil-, pyrimethanil- or mepanipyrim-containing products must not exceed the maxima as mentioned below.

For sound resistance management, good agricultural practices, including phytosanitary measures and crop protection, should be followed not only in commercial practice, but also in nurseries.

The use recommendations were reviewed during the meeting on January 17<sup>th</sup>, 2022. The *Botrytis* and *Venturia* guidelines have not been changed.

For all other pathogens and crops, specific guidelines are not yet regarded to be needed. However, the maximum number of applications should be limited and should not exceed more than 50% of all treatments.

#### **2.1.1 AP - *Botrytis* Guidelines**

- Where up to three treatments are made per season, the number of applications of AP-containing products is limited to one.
- In situations where four to six *Botrytis* treatments are made per crop and season, a maximum of two applications with AP-containing products are recommended.
- In specific situations where seven or more *Botrytis* treatments are required per crop and season, a maximum of three applications with AP-containing products is recommended and not more than two consecutive applications.
- For specific crops and products, follow use recommendations of individual companies.

#### **2.1.2 AP - *Venturia* Guidelines**

- Apply a maximum of four AP-containing products per season.
- In locations where moderately adapted or resistant populations have been reported, use APs only in mixture with an effective non-cross-resistant scab fungicide.
- Individual products should always be used at recommended dose rates and during the period when they are most effective.
- Curative use only in conjunction with reliable scab warning systems.

The next AP FRAC Working Group meeting is scheduled for January 16<sup>th</sup>, 2023.

## Literature

Mosbach *et al.*, 2017: Anilinopyrimidine Resistance in *Botrytis cinerea* is linked to Mitochondrial Function, Front. Microbiol., <https://doi.org/10.3389/fmicb.2017.02361>