



## Minutes of the FRAC OSBPI Working Group Meeting

29 September 2020 – 9:00 to 13:00

Syngenta Office, Basel (Switzerland)

### Participants

---

|          |                        |
|----------|------------------------|
| Corteva  | Jean-Luc Genet (Chair) |
|          | Olivier Couery         |
|          | Mamadou Mboup          |
| Syngenta | Helge Sierotzki        |
|          | Stefano Torriani       |
|          | Renu Kapill            |
|          | Daniela Pfeifer        |
| Bayer    | Daniela Portz          |
|          | Andreas Mehl           |
|          | Christian Zupanc       |

## Agenda

---

1. Welcome
2. Antitrust Guidelines
3. FRAC OSBPI Membership
4. Review of sensitivity monitoring data
  1. *Plasmopara viticola*
  2. *Phytophthora infestans*
  3. *Pseudoperonospora cubensis*
  4. Other pathogens
5. Review of use recommendations
6. Review of target site mutations identified so far
7. Monitoring methods
8. Next meeting

## Membership

The working group is comprised of the following members:

|                        |          |
|------------------------|----------|
| Jean-Luc Genet (chair) | Corteva  |
| Amandine Picard        | Corteva  |
| Olivier Couery         | Corteva  |
| Helge Sierotzki        | Syngenta |
| Stefano Torriani       | Syngenta |
| Jennifer Foster        | Syngenta |
| Daniela Pfeifer        | Syngenta |

## **Disclaimer**

The technical information contained in the global guidelines/the website/the publication/the minutes is provided to CropLife International/RAC members, non-members, the scientific community and a broader public audience.

While CropLife International and the RACs make every effort to present accurate and reliable information in the guidelines, CropLife International and the RACs do not guarantee the accuracy, completeness, efficacy, timeliness, or correct sequencing of such information. CropLife International and the RACs assume no responsibility for consequences resulting from the use of their information, or in any respect for the content of such information, including but not limited to errors or omissions, the accuracy or reasonableness of factual or scientific assumptions, studies or conclusions.

Inclusion of active ingredients and products on the RAC Code Lists is based on scientific evaluation of their modes of action; it does not provide any kind of testimonial for the use of a product or a judgment on efficacy. CropLife International and the RACs are not responsible for, and expressly disclaim all liability for, damages of any kind arising out of use, reference to, or reliance on information provided in the guidelines.

Listing of chemical classes or modes of action in any of the CropLife International/RAC recommendations must not be interpreted as approval for use of a compound in a given country. Prior to implementation, each user must determine the current registration status in the country of use and strictly adhere to the uses and instructions approved in that country.

## Introduction

A FRAC OSBPI Working Group was formed in 2015 to generate common resistance management recommendations for the fungicides oxathiapiprolin and fluoxapiprolin. OSBP fungicides are active against oomycete fungi and used for the control of Phytophthora and downy mildews of numerous crops.

OSBPIs inhibit an oxysterol binding protein (OSBP) homologue. Oxysterol binding proteins are implicated in the movement of lipids between membranes, among other processes. Inhibiting OSBP may disrupt other processes in the fungal cell, such as signaling, maintaining cell membranes, and the formation of more complex lipids that are essential for the cell to survive.

Oxathiapiprolin and fluoxapiprolin are cross-resistant.

OSBPIs have been classified under the FRAC Code 49. The resistance risk is medium to high.

| FRAC Code | Target site and code                         | Group name   | Chemical group                    | Common name                           | Comments  |
|-----------|--|--|-----------------------------------|---------------------------------------|---|
| 49        | F9<br>lipid homeostasis and transfer/storage | OSBPI oxysterol binding protein homologue inhibition | piperidinyl-thiazole-isoxazolines | Oxathiapiprolin<br><br>Fluoxapiprolin | Resistance risk assumed to be medium to high (single site inhibitor).<br>Resistance management required.<br>. |

Anti-Trust Guidelines (from FRAC Constitution) were shown before meeting started.

## **OSBPI Minutes of the 2020 discussions**

### **Review of sensitivity monitoring data from 2019**

#### **Grape downy mildew (*Plasmopara viticola*)**

Data presented by Bayer, Corteva and Syngenta

In 2019, sensitivity data have been generated for samples originating from Austria, Bulgaria, Croatia, France, Germany, Greece, Hungary, India, Italy, Portugal, Slovenia, Spain and Switzerland.

Most 2019 samples in monitored areas were sensitive.

Less sensitive isolates were detected at low frequency in a small number of locations in Italy (Veneto, Emilia-Romagna, Puglia), Spain (Galicia) and Germany (Baden-Württemberg).

In previous years, a few isolates with reduced sensitivity have been found in a small number of trial sites located in Austria, France, Italy, Germany, Portugal and Spain where OSBPI fungicides have been used intensively during several years. The frequency of these isolates was however low. In some of these isolates, target site mutations have been identified at positions 770, 837 and 863. More work is needed to understand the relevance of these mutations on OSBPI sensitivity.

#### **Potato/tomato late blight (*Phytophthora infestans*)**

Data presented by Bayer, Corteva and Syngenta

In 2019, sensitivity data have been generated for samples originating from Belgium, China, Colombia, Croatia, Denmark, Finland, Germany, Greece, Hungary, India, Indonesia, Ireland, Italy, Latvia, Mexico, Netherlands, Poland, Portugal, Slovakia, Spain, Sweden, Switzerland and the UK.

Most 2019 samples in monitored areas were sensitive.

Single isolates with reduced sensitivity were detected in four commercial areas of Indonesia and in the Antioquia region of Colombia.

In most of the isolates from Indonesia, a mutation has been detected at position 770.

### **Cucurbit downy mildew (*Pseudoperonospora cubensis*)**

Data presented by Corteva and Syngenta

In 2019, sensitivity data have been generated for samples originating from cucumber and melon crops in China, France, Greece, Italy, Mexico and Spain

The vast majority of the samples were sensitive.

Two populations with slightly higher EC50 values were detected in the Shandong province of China (crops grown under polytunnel). Practical relevance of this observation needs further investigations.

In past years, molecular characterization of isolates has revealed the presence of target site mutations at *Phytophthora capsici* homolog positions 700, 767 and 793.

### **Lettuce downy mildew (*Bremia lactucae*)**

Data presented by Syngenta

In 2019, sensitivity data have been generated for a limited number of samples originating from Belgium, Greece, Italy, Portugal, Poland and Spain. All the samples analyzed were sensitive.

### **Sunflower downy mildew (*Plasmopara halstedii*)**

Data presented by Corteva

In 2019, sensitivity data has been generated for samples from France, Hungary, Italy and Romania. All isolates were fully sensitive.

## **Review of global guidelines**

### **OSBPI – General Use Recommendations**

- Fungicide programs must deliver effective disease management. Apply OSBPIs at effective rates and intervals according to manufacturers' recommendations. Effective disease management throughout the season is a critical component to delay the build-up and spread of resistant pathogen populations.
- Apply OSBPIs only preventatively and in mixtures with effective fungicides from different cross-resistance groups
- The mixture partner should give effective control of the target disease(s) at the rate and interval selected.
- Foliar exposure to OSBPI products should not exceed thirty-three percent (33%) of the total period of protection needed per crop.
- The number of foliar applications of OSBPI products within a total disease management program must be limited as follows:

### **OSBPI – Grapes**

- Make no more than two (2) applications per season.

### **OSBPI – All other crops**

- Make no more than four (4) applications or maximum 33% of the total period of protection needed per crop, whichever is more restrictive.
- Where the total number of fungicide applications targeting oomycetes is less than three (3), apply no more than one (1) application of an OSBPI product.
- There should be no more than two (2) foliar applications of any OSBPI product for the control of soil-borne pathogens.
- Applications of OSBPI-containing products are to be made no more than three (3) times in sequence before applying a fungicide with a different mode of action. In areas where the agronomic risk is very high (e.g. continuous potato or cucurbit cropping) and resistance has already been reported, further restrictions to the number of consecutive applications are recommended.
- Applications of OSBPI products can be made in alternation with a fungicide with a different mode of action.

### **OSBPI – Seed/soil treatments**

- No foliar fungicide application of an OSBPI fungicide should be made following a seed/soil treatment\* with OSBPI fungicides targeting the same pathogen.

\* Directed stem sprays are interpreted as foliar not soil application.

### **OSBPI – Multiple crops**

- In case of multiple crops, do not make more than six (6) foliar applications of OSBPI product per year on the same acreage, targeting the same pathogen.
- Do not make more than one seed/soil treatment application of OSBPI per year on the same acreage, targeting the same pathogen.

### **OSBPI – Nursery crops**

- OSBPI products must not be used in nursery production of transplanted agricultural crops.

### **Next meeting:**

13 April 2021 at the Syngenta office in Basel