



STEROL BIOSYNTHESIS INHIBITOR (SBI) WORKING GROUP

Minutes from WG meeting on March 27th, and September 9th, 2025

Protocol of the discussions and recommendations of the SBI working group of the Fungicide Resistance Action Committee (FRAC)

Participants of the SBI WG Meetings

ADAMA	Dorin Pop
BASF	Martin Semar Gerd Stammler Tobias Erven (excused)
Bayer CropScience	Frank Goehlich (excused) Andreas Mehl Juergen Derpmann
Corteva Agriscience	Mamadou Kane Mboup Andrew Leader Amandine Picard
FMC	Gaelle Huet Henry Ngugi Luis Demant (excused)
Sumitomo	Yuichi Matsuzaki
Syngenta	Irina Metaeva (excused) Lorenzo Borghi Paolo Galli (excused)
UPL	Ronaldo Rodrigues (excused)

Venue of the January meeting:

Frankfurt, Hotel Westin

Hosting organization:

FRAC/Crop Life International

Disclaimer

The technical information contained in the global guidelines/the website/the publication/the minutes is provided to CropLife International/RAC members, non-members, the scientific community, and a broader public audience.

While CropLife International and the RACs make every effort to present accurate and reliable information in the guidelines, CropLife International and the RACs do not guarantee the accuracy, completeness, efficacy, timeliness, or correct sequencing of such information. CropLife International and the RACs assume no responsibility for consequences resulting from the use of their information, or in any respect for the content of such information, including but not limited to errors or omissions, the accuracy or reasonableness of factual or scientific assumptions, studies or conclusions.

Inclusion of active ingredients and products on the RAC Code Lists is based on scientific evaluation of their modes of action; it does not provide any kind of testimonial for the use of a product or a judgment on efficacy. CropLife International and the RACs are not responsible for, and expressly disclaim all liability for, damages of any kind arising out of use, reference to, or reliance on information provided in the guidelines.

Listing of chemical classes or modes of action in any of the CropLife International/RAC recommendations must not be interpreted as approval for use of a compound in a given country. Prior to implementation, each user must determine the current registration status in the country of use and strictly adhere to the uses and instructions approved in that country.

Anti-Trust Guidelines (from FRAC Constitution) were shown before meetings started.

1. DMIS AND AMINES: CEREALS

1.1. WHEAT

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

1.1.1. Septoria Leaf Blotch (*Zymoseptoria tritici* / *Mycosphaerella graminicola*)

Presentation of monitoring data for **DMIs**: ADAMA, BASF, Bayer, Corteva, FMC, Sumitomo, Syngenta

- In 2024, monitoring was carried out in Austria, Belgium, Bulgaria, Croatia, Czech Republic, Denmark, Estonia, France, Germany, Hungary, Ireland, Italy, Latvia, Lithuania, Netherlands, Norway, Poland, Romania, Slovakia, Spain, Sweden, Switzerland, Ukraine, and United Kingdom.
- Overall, the sensitivity of European populations monitored in 2024 stayed in the range observed in previous years. In general, the field performance of DMI-containing fungicides was good when used according to the manufacturers and FRAC recommendations.

In regions with limited options in fungicides classes and/or a common practice of significantly reduced rates DMIs are at higher risk and performance might be impacted.

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical data:

- In 2023, monitoring was carried out in Austria, Belgium, Bulgaria, Croatia, Czech Republic, Denmark, Estonia, France, Germany, Hungary, Ireland, Italy, Latvia, Lithuania, Netherlands, Norway, Poland, Romania, Russia, Slovakia, Spain, Sweden, Switzerland, Ukraine, and United Kingdom.
- During 2020 to 2022, monitoring was carried out in Austria, Belgium, Bulgaria, Croatia, Czech Republic, Denmark, Estonia, France, Germany, Hungary, Ireland, Italy, Latvia, Lithuania, Netherlands, Norway, Poland, Romania, Russia, Slovakia, Spain, Sweden, Switzerland, Türkiye, Ukraine, and the United Kingdom. Most of the above listed countries were covered by the monitoring programs every year, but not all.
Already in 2020, the sensitivity of populations was overall stable on the European level with EC₅₀ sensitivity values being in the range of previous years, and as in 2019, DMI EC₅₀ values were somewhat higher in the UK and Ireland than observed on the European continent where a gradient can be observed from North-West to South-East.

- In 2019, the sensitivity of the populations was overall stable on the European level with EC₅₀ sensitivity values slightly higher compared to 2018 in some geographies but overall, in the range of previous years.
- In 2018, the sensitivity of the populations was overall stable on the European level.
- In 2016 and 2017, the sensitivity of populations was overall stable on a European level with regional differences also based on different disease epidemics. In regions with lower sensitivity in 2015, the sensitivity of the populations was stable and, in some areas, even partially increased.
- In 2015, depending on the individual active ingredient and regions slight shifts of sensitivity of populations have been observed. Highest EC₅₀ values were observed in areas of elevated disease pressure and sub-optimal use of azoles in spray programs (e.g., reduction of rates in comparison to the manufacturer's recommended rate and inappropriate use of effective mix-partners).
- After the slight increase in the frequency of less sensitive isolates from 2002 to 2004, the situation had stabilised between 2005 and 2008. In 2009, a trend to slightly higher EC₅₀ values were observed in important cereal growing areas (France, Germany, Ireland, United Kingdom); this trend has slowed down in 2010 to 2012 and was stable in 2013. 2014 sensitivity was in the same range as 2011.

In *Z. tritici*, different DMI haplotypes can lead to varying levels of sensitivity depending on the chemical structure. As DMIs are generally cross-resistant, resistance management approaches should be the same for all DMIs.

1.1.2. Powdery mildew (*Blumeria graminis* f.sp. *tritici*)

Presentation of monitoring data for **DMIs**: Bayer, Corteva, Sumitomo

- In 2024, monitoring was carried out in Denmark, France, Germany, Sweden and the United Kingdom.
- Sensitivity data presented for 2016 to 2024 confirmed that the situation was overall stable within the range of variability detected during more than 20 years.

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical data:

- During 2019 to 2023, monitoring was carried out in Belgium, Czech Republic, Denmark, France, Germany, Hungary, Italy, Latvia, Lithuania, Poland, Spain, and United Kingdom. Most of the listed countries were covered by the monitoring programs every year, but not all.
- Differences in the sensitivity are significantly a.i. and regionally dependent. Higher resistance factors were observed only for particular DMIs, especially in France, Germany, and UK, but also to a lesser extent in Belgium.
- A limited monitoring in New Zealand in 2019 showed sensitivity ranges comparable to European populations.

Presentation of monitoring data for **Amines**: Bayer

- Field performance of amine-based products was good.
- In 2024, monitoring was carried out in Denmark, France, Germany, Sweden and the United Kingdom.
- Sensitivity data presented confirmed that the situation in 2024 was stable remaining in the range of variability seen over more than 25 years.

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical data:

- In 2020 and 2022, monitoring was carried out in Czech Republic, Denmark, France, Germany, Italy, Lithuania, Poland, Slovakia, and the United Kingdom. Most of the listed countries were covered by each monitoring program, but not all.

1.1.3. Wheat brown rust (*Puccinia triticina*)

Presentation of monitoring data for **DMIs**: BASF, Bayer, Syngenta, Sumitomo

- In 2024, monitoring was carried out in Denmark, France, Germany, Hungary, Ireland, the Netherlands, Poland, Sweden and the United Kingdom.
- Data from 2024 for wheat brown rust showed that sensitivities were in the range of those of more than 20 years, with a tendency to a more sensitive situation.

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical data:

- In 2020 and 2023, monitoring was carried out in Belgium, Czech Republic, Denmark, France, Germany, Hungary, Italy, Lithuania, Poland, Romania, Slovakia, Spain, and the United Kingdom. Most of the listed countries were covered by each monitoring program, but not all.

1.1.4. Eyespot (*Oculimacula* spp.)

Presentation of monitoring data for **DMIs**: Bayer, Syngenta

- In 2024, monitoring is carried out in Germany, France Lithuania and the United Kingdom
- A comparable range of sensitivity as in previous years was observed in all countries.

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical data:

- In 2023, monitoring is carried out in Germany, France and Poland.
- An analysis of samples from France, Germany, Latvia, Lithuania, Poland, Russia, Spain, Ukraine, and United Kingdom from 2020 was presented, and the same range of sensitivity observed in all countries as in previous years.
- In 2021, monitoring was carried out in Germany, Italy, Latvia, Poland, Slovakia and Ukraine.
- In 2019, still comparable sensitivity ranges and medians were observed in all monitored countries without any geographical variations.
- The 2018 data showed a homogenous and sensitive situation in all countries.
- Between 2003 and 2012 there was no change in the sensitivity of W and R types, and a stable situation had been observed during that time. In 2013, some sensitivity change has been observed in the United Kingdom, but not in France or Germany. In 2014 further sensitivity decrease has been observed in the United Kingdom, and for the first time also in France and Germany. However, overall, resistance factors still remain low, and performance was not affected.

1.1.5. Tan spot (*Pyrenophora tritici-repentis*)

Presentation of monitoring data: Syngenta

- From 2019 to 2021 and 2024, a limited monitoring was carried out in countries like Bulgaria, Czech Republic, Finland, Hungary, Lithuania, Romania, Slovakia, Sweden, and the United Kingdom.
- In these three years of monitoring, a stable and sensitive situation was observed.

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

1.1.6. Yellow rust (*Puccinia striiformis*)

Presentation of monitoring data for **DMIs**: BASF, Bayer, Sumitomo, Syngenta

- In 2024, monitoring was carried out in Denmark, France, Germany, Poland, the Netherlands and the United Kingdom.
- The first monitoring in 2015 showed high sensitivity and low diversity, and from 2016 to 2024 a stable situation was reported.

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical data:

- In 2021 to 2023, monitoring was carried out in Belgium, Czech Republic, France, Germany, Italy, Netherlands, Poland, Romania, Spain and the United Kingdom. Most of the listed countries were covered by the monitoring programs each year, but not all.

1.1.7. Snow Mould (*Microdochium nivale* & *M. majus*)

Presentation of monitoring data for **DMIs**: Bayer, Syngenta

- In 2024, only limited monitoring was carried out in Germany and Italy, .
- In general, a stable sensitivity situation has been reported for the past ten years.

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical data:

- During 2019 to 2023, monitoring was carried out in Belgium, Bulgaria, Czech Republic, France, Germany, Hungary, Italy, Romania, Russia, Spain, Sweden, Ukraine, and the United Kingdom. Most of the listed countries were covered by the monitoring programs each year, but not all.

1.1.8. Fusarium Head Blight (*Fusarium graminearum*, *F. culmorum*)

Presentation of monitoring data for **DMIs**: Bayer, Sumitomo, Syngenta

- In 2024, monitoring was carried out in Belgium, France, Germany, Hungary, the Netherlands, Poland and the United Kingdom.
- For the past 10 years, a stable sensitivity situation was observed.
- As known from previous research work different intrinsic activity was reported for other Fusarium species such as *F. poae*, *F. tricinctum*, *F. kyushuense*, as well as *F. equiseti*.

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical data:

- In 2021 and 2022, monitoring was carried out in France, United Kingdom, and Germany.
- In 2019, monitoring was carried out in France.

1.1.9. Glume blotch (*Parastagonospora nodorum*)

Presentation of monitoring data for **DMIs**: Syngenta

- No monitoring was conducted in 2024.
- In 2020 & 2021, a limited monitoring was carried out in countries like Czech Republic, Germany, Hungary, Latvia, and Sweden.
- A very narrow sensitivity range with high sensitivity levels was observed in both years

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

1.1.10. Loose wheat smut (*Ustilago tritici* / *U. nuda*)

Presentation of monitoring data for **DMIs**: Syngenta

- In 2024, a limited monitoring was carried out in countries like Austria, Bulgaria, France, Germany, Hungary, Lithuania and Poland.
- A high level of sensitivity and a narrow range of sensitivity was observed in all years.

Historical data:

- In 2020 & 2021, a limited monitoring was carried out in countries like Bulgaria, Germany, Poland, Spain, and Sweden,

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

1.2. BARLEY

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

1.2.1. Powdery Mildew (*Blumeria graminis* f.sp. *hordei*)

Presentation of monitoring data for **DMIs**: Bayer

- In 2024, monitoring was carried out in France, Germany and the United Kingdom.
- DMI products performed well. The sensitivity of the populations stayed in the range observed for almost 20 years.

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical data:

- Between 2016 and 2023, monitoring was carried out in Czech Republic, Denmark, Estonia, Germany, France, Germany, Hungary, Italy, Latvia, Poland, Sweden, Ukraine, and the United Kingdom. Country origin of samples

was varying from year to year, and monitoring was most regularly carried out in France, Germany, and the United Kingdom.

Presentation of monitoring data for **Amines**: Bayer

In 2024, monitoring was carried out in France, Germany and the United Kingdom.

The sensitivity was similar to previous years.

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical data:

- In 2020 and 2023, monitoring was carried out in Austria, Czech Republic, Estonia, France, Germany, Hungary, Italy, and United Kingdom. Most of the listed countries were covered by the monitoring programs each year, but not all.

1.2.2. Scald (*Rhynchosporium secalis*)

Presentation of monitoring data for **DMIs**: BASF, Syngenta

In 2024, monitoring was carried out in Czech Republic, France, Germany, , Hungary, Ireland, Slovakia, Romania, Spain and the United Kingdom.

- Stable situation. The sensitivity of the populations stayed in the range as observed in Europe for more than 20 years.

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical data:

- During 2020 and 2023, monitoring was carried out in Czech Republic, Denmark, France, Germany, Hungary, Ireland, Italy, Latvia, the Netherlands, Poland, Slovakia, Spain, and the United Kingdom. Most of the listed countries were covered by the monitoring programs each year, but not all.

1.2.3. Net blotch (*Pyrenophora teres*)

Presentation of monitoring data for **DMIs**: Bayer, Syngenta

- In 2024, monitoring was carried out in Austria, Belgium, Czech Republic, Denmark, France, Germany, Hungary, Ireland, Italy, Latvia, Lithuania,

Netherlands, Poland, Romania, Spain, Sweden, Ukraine and the United Kingdom.

- Variable findings across different studies were reported. Further investigations are ongoing.

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical data:

- During 2020 and 2023, monitoring was carried out in Austria, Belgium, Bulgaria, Czech Republic, Denmark, Finland, France, Germany, Greece, Hungary, Ireland, Italy, Latvia, Lithuania, Poland, Romania, Russia, Slovakia, Spain, Sweden, Switzerland, the Netherlands, Ukraine, and the United Kingdom. Most of the listed countries were covered by the monitoring programs each year, but not all. Overall, the sensitivity of populations showed no major geographical differences across Europe.
- In 2019, like 2017, lower sensitivities have been frequently detected in major French regions and in a single location in North-Eastern Germany. In the other European regions monitored sensitivity ranges were stable.
- The monitoring of the last 20 years showed a certain level of fluctuations of the sensitivity level in the regions over the years. In 2018, the situation stabilized again in all countries including France and Germany, thus being comparable to the long-term monitoring results.
- In 2017 in France significant shifts of sensitivity of populations have been observed. Highest EC₅₀ values were observed in areas of elevated disease pressure, often coupled with a reported reduced variety-resistance at significant cultivation areas, and sub-optimal use of azoles in spray programs (e.g., reduction of rates in comparison to the manufacturer's recommended rate and inappropriate use of effective mix-partners).
- In general, over the past years a significant fluctuation in sensitivity levels between the years was detected. In 2017 in single locations in Germany there have been seen some shifting which needs to be observed in the next season. The monitoring in the other countries showed a stable situation in 2017 within the regular fluctuation.

1.2.4. *Ramularia* leaf spot (*Ramularia collo-cygni*)

Presentation of monitoring data for **DMIs**: BASF, Syngenta

- In 2024, monitoring was carried out in Austria, Czech Republic, Bulgaria, France, Germany, Hungary, Ireland, Italy, Lithuania, Poland, Spain, Sweden,

Switzerland, and the United Kingdom. The results from bioassay and molecular analysis focusing on the most relevant mutations are:

- no resistance detected in single samples from Bulgaria and Italy
 - low frequencies of resistance in Spain
 - moderate frequencies of resistance in Switzerland
 - high frequencies of resistance in Austria, Czech Republic, Germany, France, Ireland, , and United Kingdom as well as in single samples originating from Hungary, Lithuania, Poland and Sweden
- Isolates were detected showing significant loss of sensitivity. Relevant CYP51-mutations explaining the effects have been identified (I325T, I328L, Y403C/Y405H).

Despite the high frequency of resistance in several countries, DMIs continue to contribute to disease control and remain an important tool for resistance management.

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical data:

- In 2023, monitoring was carried out in Croatia, Denmark, France, Germany, Hungary, Ireland, Italy, Poland, Romania, Spain, Slovenia, Switzerland, and the United Kingdom. The results from bioassay and molecular analysis focusing on the most relevant mutations are:
 - no resistance in Italy
 - low frequencies of resistance in Spain
 - moderate frequencies of resistance in Hungary, Croatia, Slovenia (variable), Denmark
 - high frequencies of resistance in Austria, Czech Republic, Switzerland, Germany, France, Ireland, Poland, Romania (limited data), and United Kingdom.
- In 2022, monitoring was carried out in Austria, Czech Republic, France, Germany, Ireland, Italy, Netherlands, Poland, Spain, Sweden, Switzerland, and the United Kingdom. The results from bioassay and molecular analysis focusing on the most relevant mutations are:
 - no resistance in Italy
 - low to moderate frequencies of resistance in Spain
 - moderate to high frequencies of resistance in The Netherlands
 - high frequencies of resistance in Austria, Czech Republic, France, Germany, Ireland, Switzerland, Poland, Sweden, Switzerland, and the United Kingdom.
- In 2021, monitoring was carried out in Austria, Croatia, Czech Republic, Denmark, France, Germany, Ireland, Italy, Netherlands, Spain, Sweden, and

United Kingdom. The results from bioassay and molecular analysis focusing on the most relevant mutations are:

- no resistance in Italy
- low frequencies of resistance in Spain and Croatia.
- moderate frequencies of resistance in Austria and The Netherlands.
- moderate to high frequencies of resistance in Czech Republic, France, Germany, Ireland, Sweden, and the United Kingdom.
- high frequencies of resistance in Denmark.

On the European continent, a gradient in terms resistance frequencies can be observed from north to south. Overall, the frequency of relevant CYP51-mutations was comparable to 2020. The field performance of DMI-containing products remains still relatively good in 2021.

- In 2020, monitoring was carried out in Denmark France, Germany, Hungary, Ireland, Italy, Lithuania, Poland, Slovakia, Spain, Sweden, Switzerland, and United Kingdom. The results from bioassay and molecular analysis focusing on the most relevant mutations are:

- no resistance in Italy
- no to low frequencies of resistance in Italy, Switzerland, and Spain.
- no to high frequencies of resistance in France.
- moderate frequencies of resistance in Austria and The Netherlands.
- moderate to high frequencies of resistance in Germany and Sweden.
- high frequencies of resistance in Czech Republic, Denmark, France, Hungary, Ireland, Lithuania, Slovakia, and United Kingdom.

- In 2019, the results are:

- no isolates/samples with the above-mentioned mutations were detected in Spain and Italy.
- no to low frequencies of resistance in Slovenia and Croatia.
- low frequencies of DMI resistance alleles were detected in Switzerland and Slovakia.
- in Austria, low to moderate frequencies were observed.
- moderate to high frequencies of resistance in Belgium, Germany, and Sweden.
- high frequencies of resistance in France, Ireland, and the United Kingdom.

- In 2018, the results are:

- no isolates with the above-mentioned mutations were detected in Italy, Spain, Sweden, and Switzerland.
- no to high frequencies of resistance in Denmark.
- low to moderate frequencies of resistance in single samples from Austria, France, and Hungary.
- low to high frequency of resistance in Germany.
- moderate to high frequencies of resistance in Belgium, Ireland, Latvia, Netherlands, and the United Kingdom.

- Data from 2017 showed high frequency of resistant strains in Denmark, Ireland, and the United Kingdom, moderate frequency in Estonia, low to moderate frequency in Sweden, and no resistant strains in Finland.
- In 2016, a broad sensitivity range has been identified, with very high frequency of highly resistant strains in southern Germany, with moderate frequency in Denmark, Ireland, Belgium, Northwestern Germany, and with low frequency in France, Austria, Sweden, and the United Kingdom. No detection of resistance in Estonia.

1.2.5. *Puccinia hordei*

Presentation of monitoring data for **DMIs**: Bayer, Syngenta

- In 2024, monitoring was carried out in Denmark, France, Germany, Sweden, and the United Kingdom, in 2023 in Denmark, France, Germany, Sweden, and the United Kingdom, in 2021 in France, Germany and Poland, as well as in 2019, 2018 and 2014 in Denmark, France, Germany, Sweden, and in the United Kingdom.
- In the last decade, a very stable situation with a narrow range of sensitivity was observed.

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

1.2.6. Smut diseases (*Ustilago* spp.)

Presentation of monitoring data for **DMIs**: Syngenta

- No monitoring was carried out in 2024.
- In 2023, monitoring was carried out in Austria, Bulgaria, Czech Republic, France, Germany, Hungary, Italy, Lithuania, Poland, Romania, and the United Kingdom.
- A very stable situation with a narrow range of sensitivity was observed for six years, with exception of a few UK isolates from 2018.

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical data:

- During 2020 and 2022, monitoring was carried out in Austria, Belgium, Bulgaria, Czech Republic, Denmark, France, Germany, Hungary, Ireland, Italy, Latvia, Lithuania, Netherlands, Poland, Romania, Slovakia, Spain,

Sweden, Switzerland, and the United Kingdom. Most of the listed countries were covered by the monitoring programs each year, but not all.

- In 2018, from specific locations in the United Kingdom product performance issues were reported. Monitoring results from the UK from 2018 revealed a number of strains with higher EC₅₀ values. All UK samples analysed in 2019 and 2020 were in a range of sensitivity comparable to 2016 & 2017. It should be considered that particular dry and warm climate conditions might negatively affect the performance of products in 2018, favouring the disease development in terms of speed and severity.

1.2.7. Barley stripe disease (*Pyrenophora graminea*)

Presentation of monitoring data for **DMIs**: Syngenta

- In 2024, few isolates from Lithuania and Sweden were tested as sensitive.
- The sensitivity range was comparable to previous years.

Please refer for the crop specific recommendations for cereals in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical data:

- From 2019 to 2023, a limited monitoring was carried out in Finland, Germany, Hungary, Poland, United Kingdom, and Sweden.

2. DMIS AND AMINES: INDUSTRIAL CROPS

2.1. SOYBEAN

Please refer for the general recommendations for use and the recommendations for Asian soybean rust to section 5. at the end of the minutes ([link](#)).

2.1.1. Asian soybean rust (*Phakopsora pachyrhizi*)

Presentation of monitoring data for **DMIs**: BASF, Bayer, Corteva, FMC, Sumitomo, Syngenta

- A sensitivity baseline has been established in Brazil based on 2005/6 data. Extensive monitoring was carried out since 2007/8 across the country.
- In 2024/2025, monitoring data from Brazil showed still a stable sensitivity situation similar to previous seasons after a slight shift observed several years ago as reported below.

Please refer for the recommendations for Asian soybean rust in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical data:

- In 2023/2024, the monitoring in Brazil showed in general a stable sensitivity situation similar to previous seasons after a slight shift observed several years ago as reported below.
- In 2022/2023, the monitoring in Brazil showed in general a stable sensitivity situation similar to previous years. In some Brazilian regions, few samples showed slightly lower sensitivity.
- In 2021/2022, the monitoring in Bolivia, Brazil and Paraguay showed in general a stable sensitivity situation similar to previous years. In some Brazilian & Paraguayan regions, single samples showed lower sensitivity.
- In 2020/21, the monitoring in Brazil, Paraguay and Bolivia showed in general a stable sensitivity situation similar to previous years. In some Brazilian regions a tendency towards lower sensitivities was observed.
- In 2019/20, across tested Brazilian and Paraguayan regions the observed sensitivity levels were on the same level as in previous years.
- In 2018/19, in some Brazilian regions a tendency towards lower sensitivities with higher variability was observed. In other regions, the sensitivity of populations was stable compared to previous years.
- In 2017/18, monitoring showed in general a stable situation as in the last years, but locally some slight shift was observable in western parts of Brazil. Despite this situation it is recognized that a regional variability in performance of DMI mixtures has been observed.
- Sensitivity shifts have been observed with a trend to stabilize in season 2010/11. This has to be seen in connection with the recommendation of an azole use in mixtures only and the introduction of a crop-free period. This trend continued in the following seasons until season 2013/14. In 2014/2015 slight shifts in sensitivity has been observed compared to 2013/14. In 2015/16 and 2016/17 the sensitivity level was on the same level as in previous years.

2.1.2. Target Spot (*Corynespora cassiicola*)

Presentation of monitoring data for **DMIs**: BASF, Bayer, FMC, Syngenta

- First studies were carried out with isolates from season 2013/14 and 2014/15 by BASF. These initial studies showed high sensitivity to DMIs.
- In season 2024/25, monitoring was carried out in Brazil. A stable sensitivity situation was still observed in comparison to previous years. Genetic analysis did not detect any target site mutation.

Historical data:

- In season 2023/24, monitoring was carried out in Brazil. A stable sensitivity situation was still observed in comparison to previous years. Genetic analysis did not detect any target site mutation.
- In season 2022/23, monitoring was carried out in Brazil. A stable sensitivity situation was observed in comparison to previous years. Genetic analysis did not detect any new or known relevant target site mutation.
- In season 2021/22, monitoring was carried out in Brazil. A stable sensitivity situation was observed in comparison to previous years.
- In season 2020/21, monitoring was carried out by BASF and Syngenta. A stable sensitivity situation was observed in comparison to previous years.
- In seasons 2019/20 and 2018/19, monitoring was carried out by BASF and Syngenta. A stable sensitivity situation was observed in comparison to previous years.
- Monitoring analysis from season 2016/17 and 2017/18 was presented by Syngenta. A stable sensitive situation was observed.
-

2.1.3. Cercospora leaf blight (*Cercospora* spp.)

Presentation of monitoring data for **DMIs**: Syngenta

- From 2022/23 to 2024/25, monitoring was carried out in Brazil. A stable sensitivity situation was observed in comparison to previous seasons.

Historical data:

- In season 2021/22, monitoring was carried out in six Brazilian regions. A stable sensitivity situation was observed in comparison to previous seasons.
- In season 2020/21, monitoring was carried out by Syngenta in Brazil. A stable sensitivity situation was observed in comparison to 2019/2020.
- In season 2019/20, an initial monitoring was carried out by Syngenta. Data showed high level of sensitivity across sampled regions in Brazil.

2.1.4. Anthracnose (*C. gloeosporioides* & *C. siamense*, involved in stem and pod rot anomalies)

Presentation of monitoring data for **DMIs**: Syngenta

- In seasons 2021/22 and 2022/23, monitoring data showed a high level of sensitivity across sampled regions in Brazil.

2.1.5. Phomopsis seed decay (*Diaporthe* spp., involved in stem and pod rot anomalies)

Presentation of monitoring data for **DMIs**: Syngenta

- In season 2021/22 and 2022/23, monitoring data showed high level of sensitivity across sampled regions in Brazil.

2.1.6. Septoria Brown Spot (*Septoria glycines*)

Presentation of monitoring data for **DMIs**: BASF

- In the years 2021 and 2022, an initial monitoring was carried out in Slovakia and Romania (2021). Data showed high level of sensitivity.

2.2. OILSEED RAPE

Please refer for the general recommendations for use to section 5. at the end of the minutes ([link](#)).

2.2.1. Phoma leaf spot and stem canker, blackleg (*Plenodomus lingam* / *Plenodomus biglobosus*)

Presentation of monitoring data for **DMIs**: BASF

- In 2022 to 2024, monitoring was carried out in Austria, Czech Republic, Denmark, France, Germany, Latvia, Lithuania, Poland, Sweden, Romania, and the United Kingdom.
- The monitoring data showed a stable sensitivity range as observed in the last 15 years.

For recommendations see General Recommendations.

Historical information:

- Between seasons 2018/19 and 2021/22, monitoring was carried out in Austria, Bulgaria, Czech Republic, Denmark, Finland, France, Germany, Hungary, Ireland, Latvia, Lithuania, Poland, Romania, Slovakia, Sweden, and the United Kingdom. Most of the listed countries were covered by the monitoring programs each year, but not all.

2.2.2. Sclerotinia stem rot, white mould (*Sclerotinia sclerotiorum*)

Presentation of monitoring data for **DMIs**: BASF, FMC, Syngenta, in previous years as well by Bayer and Corteva

- All monitoring data from 2016 to 2024 showed a stable and narrow sensitivity range with no geographical differences.

In 2024, monitoring was carried out in France, Germany and Lithuania.

Further monitoring studies are ongoing.

For recommendations see General Recommendations.

Historical information:

- Between 2016 and 2023, monitoring was carried out in Austria, Bulgaria, Czech Republic, Denmark, Estonia, France, Germany, Hungary, Latvia, Lithuania, Netherlands, Poland, Romania, Slovakia, Sweden, Ukraine, and United Kingdom. Most of the listed countries were covered by the monitoring programs each year, but not all.

2.2.3. Light leaf spot (*Pyrenopeziza brassicae*)

Presentation of monitoring data for **DMIs**: BASF

- No monitoring was conducted in 2024.
- In 2023 and 2022, monitoring was carried out with samples originating from Denmark, Ireland and United Kingdom.
- A slight shift was observed compared to highly sensitive reference strains for Ireland and United Kingdom.

Historical information:

- Monitoring programs since 2020 in Denmark or United Kingdom showed a stable sensitivity distribution.

2.3. SUGAR BEET

Please refer for the general recommendations for use to section 5. at the end of the minutes ([link](#)).

2.3.1. Leaf spot (*Cercospora beticola*)

Presentation of monitoring data for **DMIs**: BASF, Bayer, Syngenta

- In 2024, monitoring was carried out in Austria, Belgium, Croatia, Czech Republic, France, Germany, Lithuania, the Netherlands, Poland, Russia, Slovakia, Spain, Sweden, Switzerland, Ukraine and the United Kingdom.
- In 2023, monitoring was carried out in Austria, Belgium, Czech Republic, France, Germany, , Hungary, Italy, the Netherlands, Poland, Romania, Switzerland, Ukraine and the United Kingdom.
- Overall, as in previous years, European populations of *C. beticola* showed a stable sensitivity range.

Based upon the broad range of sensitivity observed in previous years, it is assumed that a shift took place before routine monitoring was set up¹².

Historical data:

- In 2022, monitoring was carried out in Austria, Croatia, Czech Republic, France, Germany, Greece, Hungary, Italy, Lithuania, Poland, Romania, Spain, Switzerland, and the United Kingdom.
- In 2020, monitoring was carried out in Belgium, Czech Republic, Denmark, France, Germany, Hungary, Italy, Lithuania, Netherlands, Poland, Romania, Spain, and Switzerland. Overall, a stable sensitivity situation was observed as in previous years.
- In 2019, monitoring was carried out in Austria, Belgium, France, Germany, Italy, Netherlands, Poland, Romania, Slovakia, Spain, and Switzerland. A stable sensitivity situation was observed as in previous years. Single isolates with increased EC₅₀ values were already detected in France and Germany in previous years but remain stable at a low frequency.
- In 2017 and 2018, monitoring was carried out in Austria, Belgium, Czech Republic, Denmark, France, Germany, Greece, Hungary, Italy, Lithuania, Netherlands, Poland, Romania, Russia, Serbia, Slovakia, Spain, Switzerland, Türkiye, Ukraine, and the United Kingdom. A stable sensitivity situation was observed as in the last 5 to 6 years. Most of the listed countries were covered by the monitoring programs in each year, but not all.

For recommendations see General Recommendations.

Presentation of monitoring data for **Amines**: BASF, Bayer, Syngenta

¹ Karaoglanidis, G.S. and Thanassouloupoulos C.C. (2002) Phenotypic instability of *Cercospora beticola* Sacc. strains expressing resistance to the sterol demethylation-inhibiting (DMI) fungicide flutriafol after cold exposure. *Journal of Phytopathology* 150, 692-696

² Karaoglanidis, G.S., Thanassouloupoulos C.C. and Ioannidis, P.M. (2001) Fitness of *Cercospora beticola* field isolates – resistant and sensitive to Demethylation Inhibitor Fungicides, *European Journal of Plant Pathology*, **107**, 337-347

- The monitoring from 2017 to 2020 revealed a stable situation with a small range of sensitivity and without geographic variations.
- No additional monitoring was done since 2020.

For recommendations see General Recommendations.

Historical data:

- Monitoring between 2017 and 2020 was carried out in Austria, Belgium, Czech Republic, Denmark, France, Germany, Greece, Hungary, Italy, Lithuania, Netherlands, Poland, Romania, Russia, Serbia, Slovakia, Spain, Switzerland, Ukraine, and the United Kingdom. Most of the listed countries were covered by the monitoring programs each year, but not all.

2.4 RICE

Please refer for the general recommendations for use to section 5. at the end of the minutes ([link](#)).

2.4.1. Narrow brown spot (*Cercospora janseana*)

Presentation of monitoring data for **DMIs**: Syngenta

- Initial sensitivity studies performed 2017 with limited number of strains indicated high and homogenous sensitivity in Indonesia.

2.4.2 *Rhizoctonia solani*

Presentation of monitoring data for **DMIs**: Syngenta

- Monitoring was carried out in China in 2015, 2016 and 2017. The monitoring indicated a stable and sensitive situation.

2.5 COTTON

Please refer for the general recommendations for use to section 5. at the end of the minutes ([link](#)).

2.5.1. Ramularia leaf blight (*Ramulariopsis gossypii*)

Presentation of monitoring data for **DMIs**: BASF, Syngenta, FMC

- In 2024, a monitoring was carried out in Brazil. The observed sensitivity range was comparable to the results from the baseline of 2011. Single isolates with higher EC₅₀ values were detected since 2018.

Presentation of monitoring data for **Amines**: BASF

- In seasons 2018 to 2020,, 2022 and 2024, monitoring was carried out in Brazil. The results showed a sensitive situation with EC₅₀ values being in a similar range over these four seasons.

2.5.2. Target spot (*Corynespora cassiicola*)

Presentation of monitoring data for **DMIs**: BASF, BAYER, FMC and Syngenta

- In 2025, monitoring was carried out in Brazil. The data showed a sensitive situation with EC₅₀ values being still in a similar range over the seven years since the monitoring started.

Historical data:

- In 2024, monitoring was carried out in Brazil. The data showed a sensitive situation with EC₅₀ values being still in a similar range over the seven years since the monitoring started.
- In 2023, monitoring was carried out in Brazil. The results showed a sensitive situation with EC₅₀ value being in a similar range over the six years since the monitoring started.
- In 2022, monitoring was carried out by BASF, BAYER, FMC, and Syngenta in Brazil. The results showed a sensitive situation with EC₅₀ value being in a similar range over the five years since the monitoring started.
- In 2019, monitoring was carried out by BASF in Brazil. The results showed a sensitive situation with EC₅₀ value being in a similar range as in 2018.
- In 2018, BASF and Syngenta performed a monitoring in Brazil. The results showed a sensitive situation with a narrow range of sensitivity.

2.6 Sunflower

Please refer for the general recommendations for use to section 5. at the end of the minutes ([link](#)).

2.6.1 Sclerotinia stem rot (*Sclerotinia sclerotiorum*)

Presentation of monitoring data for **DMIs**: Syngenta

- In 2022, monitoring was carried out in Romania and France. Data showed a narrow sensitivity range with EC₅₀ values comparable to *S. sclerotiorum* in oilseed rape and cabbage.

Historical data:

- In 2019, initial monitoring was carried in Bulgaria, Romania, and Slovakia by BASF. Data showed a narrow sensitivity range with EC₅₀ values comparable to *S. sclerotiorum* in oilseed rape.

2.6.2 Black Stem disease (*Plenodomus lindquistii*)

Presentation of monitoring data for **DMIs**: BASF

- In 2022, initial monitoring was carried out with samples from Romania. Data showed a narrow sensitivity range.

2.7 Pea

Please refer for the general recommendations for use to section 5. At the end of the minutes ([link](#)).

2.7.1. Ascochyta blight (*Ascochyta pisi*, *Didymella pinodes*, *D. pinodella*)

Presentation of monitoring data for **DMIs**: BASF

- In 2022, monitoring was carried out in France. Data showed a narrow sensitivity range.

Historical data:

- In 2021, initial monitoring was carried out in France and Spain by BASF. Data showed a narrow sensitivity range.

2.8 Beans

Please refer for the general recommendations for use to section 5. At the end of the minutes ([link](#)).

2.8.1 Sclerotinia stem rot (*Sclerotinia sclerotiorum*)

Presentation of monitoring data for **DMIs**: Syngenta

- In 2022, monitoring was carried out in Belgium, France and the Netherlands. Data showed a narrow sensitivity range with EC₅₀ values comparable to *S. sclerotiorum* in oilseed rape and cabbage.

Historical data:

- In 2020, for the first-time monitoring on beans and green beans was carried out in France, and in 2021, samples from France and Belgium were analyzed. All isolates collected showed a narrow sensitivity range and high sensitivity (low EC₅₀ values).

3. DMIS AND AMINES: OTHER CROPS

3.1. GRAPE VINE:

Please refer for the crop specific recommendations for grape vine in combination with the general recommendations for use to section 5. At the end of the minutes ([link](#)).

3.1.1 Powdery mildew (*Erysiphe necator*)

Presentation of monitoring data for **DMIs**: BASF, Bayer, Corteva, and Syngenta

In 2024, monitoring was carried out in Austria, Bulgaria, , France, Germany, Italy, Portugal and Spain.

- Sensitivity monitoring based on bioassays showed a stable and homogeneous distribution in the range of the previous year.

Exclusive frequency measurements of a single cyp51 mutation, such as Y136F, are not sufficient to describe the sensitivity situation in *Erysiphe necator* populations towards DMIs.

Please refer for the crop specific recommendations for grape vine in combination with the general recommendations for use to section 5. At the end of the minutes ([link](#)).

Historical information:

- In 2023, monitoring was carried out in Austria, Bulgaria, Croatia, Czech Republic, France, Germany, Greece, Italy, Hungary, Portugal, Spain, Slovakia, Slovenia, Romania, Switzerland, and Türkiye.
- In 2021, monitoring was carried out in Austria, Croatia, France, Germany, Greece, Italy, Hungary, Portugal, Spain, Switzerland, and Türkiye. Generally, population sensitivity can vary significantly between locations and years within individual countries. In 2021, a shift to lower sensitivities was observed in some samples. Variable resistance factors were observed.

- Between 2017 and 2020, monitoring was carried out in Austria, Bulgaria, Croatia, Czech Republic, France, Germany, Greece, Hungary, Italy, Portugal, Romania, Slovakia, Slovenia, Spain, Switzerland, and Türkiye. Most of the listed countries were covered by the monitoring programs each year, but not all.

Presentation of monitoring data for **Amines**: Bayer

- In 2024, monitoring was carried out in Austria, Hungary, Germany, France, Italy, Portugal, Spain and Switzerland.
- Stable situation in the European countries with low resistance factors towards amines with only small regional fluctuations close to the baseline.

Please refer for the crop specific recommendations for grape vine in combination with the general recommendations for use to section 5. At the end of the minutes ([link](#)).

Historical data:

- In 2022, monitoring was carried out in Austria, Hungary, Germany, France, Italy, Portugal, and Spain.
- In 2019, monitoring was carried out in Austria, Croatia, Germany, Italy, Portugal, and Spain, as well as in 2020 in Austria, France, Germany, Hungary, Italy, Spain, and Switzerland.
A stable situation in the European countries with low resistance factors towards amines was observed, with only small regional fluctuations close to the baseline.

3.2. STONE AND POME FRUIT

Please refer for the crop specific recommendations in pome- and stonefruit in combination with the general recommendations for use to section 5. At the end of the minutes ([link](#)).

3.2.1. Scab on APPLE (*Venturia inaequalis*)

Presentation of monitoring data for **DMIs**: BASF, Bayer, and Syngenta

- Monitoring for samples from season 2024 is ongoing.
- Overall, the sensitivity in European populations remains unchanged since around a decade. A few outliers with lower sensitivity levels were observed.
- In 2023, monitoring was carried out in Austria, Belgium, Switzerland, Germany, Spain, France, Greece, Croatia, Hungary, Italy, Netherlands, Poland, Portugal, Serbia, Slovenia, Türkiye, UK.

Please refer for the crop specific recommendations in pome- and stonefruit in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical information:

- In 2022, monitoring was carried out in Belgium, Bulgaria, Croatia, France, Germany, Greece, Hungary, Italy, Latvia, Lithuania, Netherlands, Poland, Portugal, Romania, Slovenia, and Spain.
- Between 2018 and 2021, monitoring was carried out in Austria, Belgium, Bulgaria, Croatia, Denmark, France, Germany, Greece, Hungary, Italy, Latvia, Netherlands, Poland, Portugal, Romania, Slovenia, Spain, Sweden, Switzerland, and Türkiye.

3.2.2. Powdery mildew (*Podosphaera leucotricha*) on APPLE

Presentation of monitoring data for **DMIs**: Syngenta, BASF

In 2024, monitoring was carried out in Austria, Belgium, France, Germany, Greece, Hungary, Italy, the Netherlands, Portugal, Poland, Spain and Türkiye.

- Monitoring was started across Europe in 2010. No change in sensitivity since then was observed.

Please refer for the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical information:

- Between 2017 and 2023, monitoring was carried out in Austria, Belgium, Bulgaria, Croatia, Czech Republic, France, Germany, Greece, Hungary, Spain, Italy, Netherlands, Türkiye, Hungary, Latvia, Lithuania, Poland,

Portugal, Romania, Slovenia, and Spain, but some of the listed countries were not covered by the monitoring programs each year.
No change in sensitivity comparing 2022 to 2010 was observed.

3.2.3. Brown rot on stonefruit (*Monilinia* spp.) – ALMOND, APRICOT, CHERRY, NECTARINE, PEACH, PLUM

Presentation of monitoring data for **DMIs**: Syngenta and BASF

- No monitoring was conducted in 2024.
- In 2023, monitoring was carried out in Belgium, Bulgaria, France, Germany, Greece, Hungary, and Spain. Crops sampled were apricots, cherry, nectarine, other *Prunus* species and pome fruits.
- Until now, a narrow and homogenous distribution of sensitivity is detected across all crops, countries, and species.

Please refer for the crop specific recommendations in pome- and stonefruit in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical information:

- In 2021, monitoring was carried out in Croatia, Czech Republic, Belgium, Bulgaria, France, Germany, Greece, Hungary, Italy, Poland, Romania, Spain. Crops sampled were apricots, cherry, nectarine, and other *Prunus* species.
- In 2020, monitoring was carried out in Greece, France, Hungary, and Spain. Crops sampled were almond, nectarine and cherry.
- In 2018 & 2019, monitoring was carried out in Belgium, Bulgaria, Croatia, France, Germany, Hungary, Italy, Poland, and Spain.

3.2.4. *Stemphylium vesicarium* on PEARS

Presentation of monitoring data for **DMIs**: Syngenta

- No monitoring was conducted in 2024.
- In 2021, monitoring was carried out in Belgium, Italy, and Portugal.
- Overall, a homogenous and stable situation in terms of sensitivity was observed.

Please refer for the crop specific recommendations in pome- and stonefruit in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical information:

- Between 2018 and 2020, monitoring was carried out in Belgium, Hungary, Italy, Netherlands, Portugal, and Spain. Most of the listed countries were covered by the monitoring programs each year, but not all.

3.3. TOMATO / POTATO

Please refer for the general recommendations for use to section 5. at the end of the minutes ([link](#)).

3.3.1. *Alternaria solani*, *Alternaria alternata* and *Alternaria linariae*

Presentation of monitoring data for **DMIs**: BASF, Bayer, Syngenta

- In 2024, monitoring was carried out in Austria, Belgium, Czech Republic, Germany, Finland, France, Netherlands, Poland, Portugal, Serbia, and Sweden..Further monitoring studies are ongoing.
- A homogenous sensitivity of all studied *Alternaria* species was observed in 2024, comparable to previous years.

Isolates with mutation-combinations (L143F + G446S or G462S) were found. Resistance factors were very low and fitness penalties were observed.

Historical information:

- Between 2019 and 2023, monitoring was carried out on potatoes and/or tomatoes in Austria, Belgium, Bulgaria, Croatia, Denmark, Finland, France, Germany, Greece, Hungary, Italy, Latvia, Netherlands, Norway, Poland, Portugal, Romania, Serbia, Spain, Sweden and Switzerland. Most of the listed countries were covered by the monitoring programs each year, but not all.

3.3.2. *Oidium neolycopersici*

Presentation of monitoring data for **DMIs**: Syngenta

- Monitoring in 2018 showed a comparable sensitivity range as monitored since 2015, with no variations between countries.

3.3.3. Leaf mold (*Fulvia fulva*)

Presentation of monitoring data for **DMIs**: Syngenta

- In 2019, initial monitoring studies were carried out in China, and indicated a high and homogenous sensitivity.

3.3.4. Silver scurf (*Helminthosporium solani*)

Presentation of monitoring data for **DMIs**: Syngenta

- In 2023, a monitoring was conducted in Belgium and single isolates were analyzed from Germany, Hungary and Italy.
- A stable sensitivity range was observed in European countries since 2012.

Historical data:

- In 2022, potato crops grown in Belgium, Germany, France, Hungary, Italy, Netherlands, Spain, and United Kingdom were monitored.
- In 2019 and 2020, a few isolates showed slightly lower sensitivity levels.

3.4. CUCURBITS

Please refer for the general recommendations for use to section 5. at the end of the minutes ([link](#)).

3.4.1. *Podosphaera xanthii*

Presentation of monitoring data for **DMIs**: BASF and Syngenta

- In 2024, monitoring was carried out in France, Greece, , Spain, and Türkiye.
- No change of sensitivity has been observed from 2011 to 2024, and no variations between countries and samples collected from cucumbers, melon, pumpkin, or zucchini were monitored.

Historical information:

- In 2022, monitoring was carried out in France, Greece, Italy, Portugal, Spain, and Türkiye.
- In 2020, monitoring was carried out in Belgium, Bulgaria, France, Hungary, Greece, Italy, Poland, Serbia, and Spain. Crops sampled were cucumber, melon, and zucchini.

- In 2018 and 2019, monitoring was carried out in Belgium, France, Greece, Italy, Netherlands, Poland, and Spain. Most of the listed countries were covered by the monitoring programs in each year, but not all.

3.4.2. Gummy Stem blight (*Stagonosporopsis cucurbitacearum*)

Presentation of monitoring data for **DMIs**: Syngenta

- Monitoring started in 2017 and continued in 2018 and 2019 in Belgium and Spain, as well as in two Spanish locations in 2021.
- In general, a stable sensitivity situation was observed. Compared to previous years, single isolates with higher EC₅₀ values were reported in 2021.

3.5. OTHER VEGETABLES

Please refer for the general recommendations for use to section 5. at the end of the minutes ([link](#)).

3.5.1. *Alternaria* species on BROCCOLI, CABBAGE, CARROTS, CAULIFLOWER

Presentation of monitoring data for **DMIs**: BASF, Syngenta

- In 2023, monitoring was carried out in Belgium and Netherlands (limited data).
- Monitored species was *Alternaria dauci* on carrots, and no indication of decreased sensitivity has been observed.

Historical data:

- In 2022, monitoring was carried out in France. Monitored species were *Alternaria alternata* on cabbage and carrots, without any indication of decreased sensitivity across all crops, countries, and species.
- In 2019, monitoring was carried out in Belgium, Croatia, France, Greece, Hungary, Italy, Poland, Portugal, and Spain. Monitored species were *Alternaria alternata*, *A. brassicae*, *A. brassicicola* and *A. dauci* on broccoli, cabbage, and carrots, without any indication of decreased sensitivity across all crops, countries, and species.

3.5.2. *Stemphylium vesicarium* on ASPARAGUS

Presentation of monitoring data for **DMIs**: Syngenta

- In 2018, a stable sensitivity as in the previous years was observed in the United Kingdom; single isolates with higher EC₅₀ values were already detected but remained stable at a low frequency.

3.5.3. *Stemphylium botryosum* on SPINACH

Presentation of monitoring data for **DMIs**: Syngenta

- Initial sensitivity studies performed in 2018 with limited number of strains indicated high and homogenous sensitivity in the USA.

3.5.4. White mould (*Sclerotinia sclerotiorum*) on LETTUCE

Presentation of monitoring data for **DMIs**: Syngenta

- In 2020, for the first-time sensitivity monitoring on lettuce was carried out in Spain. In 2021, samples from France, Italy, and Spain were analyzed.
- All isolates collected showed a narrow sensitivity range and high sensitivity. (low EC₅₀ values).

3.5.5. *Stemphylium vesicarium* on ONIONS

Presentation of monitoring data for **DMIs**: Syngenta

- In 2021, all isolates monitored from onion collected in Germany and Poland showed EC₅₀ values in a range of known sensitivity levels of *S. vesicarium* from other crops.

3.6. CITRUS, STRAWBERRY

Please refer for the general recommendations for use of DMIs to section 5. at the end of the minutes ([link](#)).

3.6.1. Anthracnose (*Colletotrichum acutatum*)

Presentation of monitoring data for **DMIs**: Syngenta

- Initial sensitivity studies performed in 2017 with limited number of strains indicated high and homogenous sensitivity in USA.

3.7. BANANA

3.7.1. Black Sigatoka (*Pseudocercospora fijiensis*)

In case you are interested in background information for resistance management in bananas, please follow this link: [Information on Banana](#)

4. KETO-REDUCTASE-INHIBITORS – KRI (SBI-CLASS III)

This group comprises of Fenhexamid and Fenpyrazamine as inhibitors of the enzyme Keto-Reductase (KRIs). Both are cross-resistant.

4.1. Grey mould (*Botrytis cinerea*) on GRAPE VINE

Presentation of monitoring data for **KRIs**: Bayer and Sumitomo

- In 2024, monitoring was carried out in France, Germany, and Italy, Spain. The frequency of resistant isolates was low in all studied countries.

Field performance of botryticides is most effective if embedded in sound spray programmes, respecting the individual resistance management recommendations.

Please refer for the disease specific recommendations for control of *Botrytis* spp. in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical data:

- In 2023, monitoring was carried out in France, Germany, and Italy, Spain. The frequency of resistant isolates was low in all studied countries.
- In 2022, monitoring was carried out in France, Germany, Italy, and Spain. The frequency of resistant isolates was low to moderate in in all studied countries, depending on the region.
- In 2021, monitoring was carried out in France, Germany, Italy, and Spain. The frequency of resistant isolates was low in Italy, and low to moderate in France, Germany, and Spain.
- In 2020, monitoring was carried out in France, Germany, Italy, and Spain. The frequency of resistant isolates was low to moderate in France, Italy, and Spain, and moderate in Germany.
- In 2019, monitoring was carried out in Austria, France, Germany, Hungary, Italy, and Spain. The frequency of resistant isolates was low in Austria, Hungary, and Spain, low to moderate in France and Germany, and moderate in Italy.
- In 2018, the frequency of resistant isolates was very low in Hungary and Italy, low in France, moderate in Germany, and moderate to high in Chile.

- In 2017, the frequency of resistant isolates was low in Austria and France, moderate in Germany, and in Italy all strains analysed were fully sensitive.
- In 2016, moderate to high frequencies in Germany, low frequencies in France and very low frequencies in Italy and Spain.
- High frequencies of resistant isolates were detected in Chile (2014, 2015, 2016).

4.2. Grey mould (*Botrytis cinerea*) on STRAWBERRIES

Presentation of monitoring data for **KRIs**: Bayer and Sumitomo

- In 2023, monitoring was carried out in France, German, Italy, Spain, Poland and the United Kingdom. A low to moderate frequency of resistant isolates was detected in France and Poland. Moderate frequency of resistant isolates was detected Italy. High frequencies were detected in Spain and the United Kingdom.

Please refer for the disease specific recommendations for control of *Botrytis* spp. in combination with the general recommendations for use to section 5. At the end of the minutes ([link](#)).

Historical data:

- In 2023, monitoring was carried out in France, Italy, Spain and Poland. A low to moderate frequency of resistant isolates was detected.
- In 2022, monitoring was carried out in France, Germany, Spain, and Italy. The frequency of resistant isolates was low to moderate in Germany and France. In Italy and Spain, a low to high frequency was detected, dependent on the region.
- In 2021, monitoring was carried out in Poland, Germany, Italy and France. The frequency of resistant isolates was low to moderate in all countries.
- In 2020, monitoring was carried out in France, Germany, Italy, Netherlands, Norway, Poland, Spain, and the United Kingdom. The frequency of resistant isolates was low to moderate in France, Germany, Norway, and Poland, moderate in Italy, Netherlands, and the United Kingdom, and moderate to high in Spain.
- In 2019, monitoring was carried out in Austria, France, Denmark, Italy, Poland, Germany, and the United Kingdom. The frequency of resistant isolates was low in Austria, France, and Denmark, low to moderate in Poland, moderate in Germany, and moderate to high in Italy and the United Kingdom.

- In 2018, monitoring was carried out in Austria, Denmark, France, Germany, Italy, Netherlands, Norway, Poland, Sweden, Spain, and the United Kingdom. The frequency of resistant isolates was low in Austria, France, Poland, and Sweden, low to moderate in Denmark, Italy, the Netherlands and Germany, moderate in Spain, and moderate to high in Norway and the United Kingdom.
- In 2017, monitoring was carried out in 2017 in Denmark, France, Germany, Netherlands, Poland, and the United Kingdom. High presence of resistant strains was observed in the United Kingdom, low to moderate presence in Denmark, Germany, and the Netherlands, and low presence in Austria, France, and Poland.

4.3. Grey mould (*Botrytis cinerea*) on RASPBERRIES

Presentation of monitoring data for **KRIs**: Bayer

- Monitoring in 2015 in the Netherlands showed moderate frequency of resistance.
- Limited monitoring in Norway in 2014 showed high frequency of resistant strains.

Please refer for the disease specific recommendations for control of *Botrytis* spp. in combination with the general recommendations for use to section 5. At the end of the minutes ([link](#)).

4.4. Grey mould (*Botrytis cinerea*) on TOMATO

Presentation of monitoring data for **KRIs**: Sumitomo

- In 2024, monitoring was carried out in Italy, and France. Frequency was low dependent on the region.

Please refer for the disease specific recommendations for control of *Botrytis* spp. in combination with the general recommendations for use to section 5. at the end of the minutes ([link](#)).

Historical data:

- In 2023, monitoring was carried out in Spain, Italy, and France. Frequency was low to moderate dependent on the region, thus decreased compared to the previous year.

- Monitoring was carried out in Spain (2020), Italy and France (2021), Spain, Italy, and France (2022).
- In previous years, monitoring was carried out in France, Hungary, Italy, the Netherlands, Poland, and Spain. The frequency of resistant isolates in Spain, Italy, and France was low. In 2022, frequency was moderate to high dependent on the region.
- In 2019, monitoring was carried out in France, Hungary, and Italy. The frequency of resistant isolates was low in France and Hungary, and moderate in Italy.

4.6. Brown rot (*Monilinia* spp.) in STONE FRUITS

Presentation of monitoring data for **KRIs**: Sumitomo

- In 2024, monitoring was carried out in Spain, France and Italy.
- A stable sensitivity range was observed as in previous years.

For recommendations see General Recommendations.

Historical information:

- In 2023, monitoring was carried out in Spain, France and Italy.
- In 2022, monitoring was carried out in Spain and Italy.
- In 2021, monitoring was carried out in Spain and Italy, as well as in 2020 in France, Italy, and Spain.

5. Review of global guidelines

5.1 SBI – General recommendations for use

The SBI fungicides represent one of the most potent classes of fungicides available to the grower for the control of many economically important pathogens. It is in the best interest of all those involved in recommending and using these fungicides that they are utilised in such a way that their effectiveness is maintained.

The working group concentrates its resources on the major crop/pathogen targets from the point of view of resistance risk. Inevitably many, still important pathogens are omitted. To help in making recommendations for crops and pathogens not directly covered, the following general recommendations can be made:

- Repeated application of SBI fungicides alone should not be used on the same crop in one season against a high-risk pathogen in areas of high disease pressure for that particular pathogen.
- For crop/pathogen situations where repeated spray applications (e.g., orchard crops/powdery mildew) are made during the season, alternation (block sprays or in sequence) or mixtures with an effective non-cross-resistant fungicide are recommended.
- Where alternation or the use of mixtures is not feasible because of a lack of effective or compatible non-cross-resistant partner fungicides, then input of SBI's should be reserved for critical parts of the season or crop growth stage.
- If the performance of SBIs should decline and sensitivity testing has confirmed the presence of less sensitive isolates, SBIs should only be used in mixture or alternation with effective non-cross-resistant partner fungicides.
- The introduction of new classes of chemistry offers opportunities for more effective resistance management. The use of different modes of action should be maximized for the most effective resistance management strategies.
- Users must adhere to the manufacturers' recommendations. In many cases, reports of "resistance" have, on investigation, been attributed to cutting recommended use rates, or to poorly timed applications.
- Fungicide input is only one aspect of crop management. Fungicide use does not replace the need for resistant crop varieties, good agronomic practice, plant hygiene/sanitation, etc.
- Exclusive frequency measurements of single cyp51 mutations are not sufficient to describe the sensitivity situation towards DMIs but can help to better understand the background of sensitivity shifts.

5.2 SBI – Recommendations for cereals (DMIs and amines)

The recommendations for the use of DMI and amine fungicides in mixture or alternation programmes with different mode of action fungicides remain unchanged.

It needs to be emphasized that it is essential for resistance management purposes to follow strictly the manufacturer's and FRAC recommendations.

- Repeated application of DMI or amine fungicides alone should not be used on the same crop in one season against risky pathogens (e.g., cereal powdery mildews, barley net blotch, scald) in areas of high disease pressure for that particular pathogen.
- Reduced rates of DMIs can contribute to accelerate the shift to less sensitive populations. It is critical to use effective rates of DMIs in order to ensure robust disease control and effective resistance management. DMIs must provide effective disease control and be used at manufacturers recommended rates.
- When used in mixture recommended effective rates of the SBI must be maintained.
- Split and reduced rate programmes, using multiple repeated applications at dose rates below manufacturer's recommendations, provide continuous selection pressure and accelerate the development of resistant populations, and therefore must not be used.

To ensure good performance and particularly resistance management in situations of even low disease pressure it is essential to adhere to dosages and spray timings as recommended by manufacturers. Curative applications should be avoided. Application timing has to be appropriate to all mix partners' characteristics. Mixing with a non-cross resistant fungicide at effective dose rates contributes to a more effective disease control and resistance management.

The amine fungicides are effective non-cross-resistant partner fungicides for DMIs on cereals for the control of pathogens included in the label recommendation of each respective product.

5.3 SBI – Recommendations for Asian soybean rust:

Refer to the general recommendations for SBI's.

In addition, to ensure robust disease control and resistance management it is essential to

- Apply DMI fungicides always in mixtures with effective non-cross resistant fungicides (mix partner shall provide control over the spraying interval).
- Refer to manufacturers recommendations for rates. Reduced rates must be avoided.
- Apply preventively or as early as possible in the disease cycle.
- Ensure a proper coverage of the treated crop by appropriate and well calibrated application technology (e.g., to ensure penetration into canopy).
- Apply DMI fungicide containing products always at intervals recommended by the manufacturers and adjusted to the disease epidemics. Avoid extended spray intervals.

- Good agricultural practices must be considered to reduce source of inoculum, disease pressure and resistance risk, e.g., no multiple cropping, implement and respect soybean-free periods, consider partially resistant soybean varieties, reduce the planting window, give preference to early-cycle varieties and endorse the destruction of volunteers.

5.4 SBI – Recommendations for Grapevine:

- DMIs and amines should be used preventative and curative situations should be avoided.
- The existing strategy for effective disease control and resistance management continues to be successful and the use recommendation is a maximum of 4 applications per season per mode of action. The strategy includes the use of mixtures or alternation with non-cross-resistant fungicides.
- To ensure that SBIs can remain the effective basis for control of *Erysiphe necator* in grapevine, their use should adhere to the full recommended rate (either alone or in mixture) at the recommended timing and application volume and an accurate treatment of each row.

5.5 SBI – Recommendations in Pome- and Stonefruit:

- DMI fungicides are not recommended for season long use and a maximum of 4 DMI sprays either alone or in mixture is recommended.
- DMIs should be used in mixtures or (block) alternations with a non-cross-resistant fungicide. Application of recommended label rates is important.
- Preventative applications should always be the first choice with DMIs. Curative applications are only recommended when accurate disease warning systems are available.

5.6 KRI – Recommendations for control of *Botrytis* spp.:

- Use KRIs only protectively.
- Use KRIs only in strict alternation, no block application
- Solo product as part of alternation programmes:
 - Spray programmes with a maximum of 3 treatments per season: max. 1 application with KRIs
 - Spray programmes with 4-5 treatments/season: max. 2 applications with KRIs
 - Spray programmes with 6 and more treatments: at the maximum one third of all Botryticide-applications
- Use in mixtures:
 - Both partners - if applied alone at the dose used in the mixture - must have sufficient activity against Botrytis.

- Not more than 50% of all Botryticide-treatments should be made with KRIs-containing mixtures.

For sound resistance management, good agricultural practices, including phytosanitary measures and crop protection, should be followed carefully.

5. NEXT MEETING

Next WG meeting is planned for March 2026.