



Minutes of the FRAC QioSI Working Group Meeting

26 March 2026, Frankfurt, Germany

Participants

Gerd Stammler (QioSI Chair)	BASF
Helge Sierotzki (QI Chair)	Syngenta
Michael Merk	BASF
Raffaello Zito	BASF
Andreas Mehl	Bayer
Chris Cooksley	Bayer
Mamadou Mboup	Corteva
Amandine Picard	Corteva
Andy Leader	Corteva
Gaelle Huet	FMC
Yoshitake Desaki	Ishihara Sangyo Kaisha
Yutaka Sakinaga	ISK Bioscience Europe
Victor de Amorn	ISK Biosciences Europe
Yasuhiro Fukami	Nissan Chemical Corporation
Hiroyuki Ito	Nissan Chemical Europe
Yuichi Matsuzaki	Sumitomo
Lorenzo Borghi	Syngenta
Stefan Dragos	Syngenta
David Ranner	Syngenta

Disclaimer

The technical information contained in the global guidelines/the website/the publication/the minutes is provided to CropLife International/RAC members, non-members, the scientific community and a broader public audience.

While CropLife International and the RACs make every effort to present accurate and reliable information in the guidelines, CropLife International and the RACs do not guarantee the accuracy, completeness, efficacy, timeliness, or correct sequencing of such information. CropLife International and the RACs assume no responsibility for consequences resulting from the use of their information, or in any respect for the content of such information, including but not limited to errors or omissions, the accuracy or reasonableness of factual or scientific assumptions, studies, or conclusions.

Inclusion of active ingredients and products on the RAC Code Lists is based on scientific evaluation of their modes of action; it does not provide any kind of testimonial for the use of a product or a judgment on efficacy. CropLife International and the RACs are not responsible for, and expressly disclaim all liability for, damages of any kind arising out of use, reference to, or reliance on information provided in the guidelines.

Listing of chemical classes or modes of action in any of the CropLife International/RAC recommendations must not be interpreted as approval for use of a compound in a given country. Prior to implementation, each user must determine the current registration status in the country of use and strictly adhere to the uses and instructions approved in that country.

1. Introduction

The FRAC QioSI (Quinone inside and outside inhibitor, stigmatellin binding mode) Working Group was formed in 2025 and the first meeting took place in March 2025. Currently, ametoctradin is the only registered fungicide with this mode of action.

QioSI fungicides control oomycete plant pathogens by inhibiting both ubiquinone binding sites at complex III (Zhu et al. 2015, Fehr et al. 2016). To date, two resistance mechanisms have been identified in *Plasmopara viticola* that affect ametoctradin sensitivity: overexpression of the alternative oxidase, which impacts all complex III inhibitors, and the S34L mutation in the cytochrome *b* gene, specific to ametoctradin. The S34L mutation occurs in the inner binding site but does not impact Qil fungicides. Resistance mechanisms for QoI (G143A and F129L) or Qils (L201S, inserts at 203/204) do not influence the sensitivity to ametoctradin.

QioSI fungicides have been classified under the FRAC Code 45. The resistance risk is assumed to be medium to high.

FRAC Code	Target site and code	Group name	Chemical group	Common name	Comments
45	C8 complex III: cytochrome bc1 (ubiq. reductase) at Qi and Qo site (stigmatellin binding mode)	QioSI fungicide (Quinone inside and outside inhibitor, stigmatellin binding mode)	triazolo-pyrimidylamine	ametoctradin	not cross-resistant to QoI fungicides, resistance risk assumed to be medium to high (single site inhibitor) Resistance Management required

The working group collects sensitivity monitoring data and recommends resistance management strategies.

References

Zhu X., Zhang M., Liu J., Ge J., Yang G. (2015) Ametoctradin is a potent Qo site inhibitor of the mitochondrial respiration complex III. *J Agric Food Chem*, 63, 3377–3386, DOI: 10.1021/acs.jafc.5b00228

Fehr M., Wolf A., Stammler G. (2016) Binding of the respiratory chain inhibitor ametoctradin to the mitochondrial bc1 complex. *Pest Manag Sci* 2016, 72, 591–602. DOI: 10.1002/ps.4031

2. Resistance monitoring

2.1. *Plasmopara viticola*

Companies: BASF, Corteva

Main resistance mechanisms are the overexpression of the alternative oxidase and the target site mutation S34L in the cytochrome *b*. Sensitivity monitoring methods include biotests (detached leaf tests, zoospore release tests) and molecular genetic methods for detection of target site mutation S34L.

In 2025 monitoring was performed in Austria, France, Germany, Greece, Hungary, Italy, Portugal Romania and Spain using biotests and molecular genetic methods (quantitative detection of the S34L mutation). Low frequencies of specific QioSI resistance were detected in Austria, France, Spain, Italy and Germany. All samples from Hungary, Greece, Portugal and Romania were sensitive.

Historical background:

In 2023 and 2024 extensive sensitivity monitoring with biotests and molecular genetic methods for S34L detection showed that specific QioSI resistance can be found at low frequency in in France, Italy and Spain. Single cases have been found in Hungary and Germany. Samples from Portugal, Greece and Türkiye showed full sensitivity.

The overexpression of AOX has been recognized in *Plasmopara viticola* for over 15 years, exhibiting regional variations. Research indicates that this mechanism is associated with fitness penalties, which may explain why AOX fluctuates in presence within vineyards and does not increase in prevalence across European viticultural regions. The target site mutation S34L has first been found in the Armagnac area in France in 2016 and afterwards in some other grape growing regions.

2.2. *Phytophthora infestans*

Companies: BASF

Sensitivity monitoring was conducted using microtiter tests with determination of EC₅₀ values.

In 2025, samples originating from Denmark, Germany, Italy, Latvia, Lithuania, Poland, and Portugal were investigated. All samples were sensitive, besides one outlier from Poland.

Historical background:

In 2024, all isolates tested were in the baseline sensitivity range and there was no indication of any adaptation. Isolates were taken from potatoes from Belgium, Greece, Italy, Latvia, Lithuania, Netherlands, Portugal, and Spain.

Data from 2019 to 2024 showed a narrow range of EC₅₀ values of *Phytophthora infestans* with an unimodal distribution. Single outliers with EC₅₀ values outside the baseline range have been identified in 2021 and 2022. A target site mutation couldn't be identified in the cytochrome b gene of such outliers. Outliers were not detected in populations analyzed in 2023.

2.3. *Pseudoperonospora cubensis*

Companies: BASF

Studies on isolates from 2025 included France, Greece, Portugal, Spain and Türkiye. Isolates from Greece were not completely inhibited and are currently further investigated. All other samples were within the baseline sensitivity.

Sensitivity monitoring was conducted using biotests with determination of EC₅₀ values. The data from 2022, 2023 and 2024 showed no indication of resistance development towards QioSI in this species. Samples were collected from Portugal, France, Italy and Spain.

3. Recommendations for Use

Resistance management strategies might differ in regions or countries because of different disease pressure or national guidelines.

Recommendations for grape downy mildew (*Plasmopara viticola*):

- Apply QioSI containing products in a preventative manner.
- Apply a maximum of 3 applications per season.
- Use always in mixture (ready-mix or tank-mix) with an effective partner
- In order to reduce selection pressure, use in alternation with a non cross-resistant product is recommended.

Recommendations for late blight (*Phytophthora infestans*)

- Apply QioSI containing products in a preventative manner.
- Apply a maximum of 50 % of the total number of intended applications for late blight control during one crop cycle.
- Mixtures (ready-mix or tank-mix) with an effective partner are recommended.
- Alternation with fungicides having other modes of action is recommended in spray programs.
- Apply QioSI fungicides according to manufacturers' instructions.

Recommendations for oomycetes occurring in vegetable crops

- Apply QioSI containing products in a preventative manner.
- Mixtures (ready-mix or tank-mix) with an effective partner are recommended.
- Alternation with fungicides having other modes of action is recommended in spray programs.
- Apply QioSI fungicides according to manufacturers' instructions.

4. Next Meeting

Next annual meeting is scheduled for 18th March 2027.